

GEOMETRIC MEASURES FOR HYPERBOLIC SETS ON SURFACES

A. A. PINTO AND D. A. RAND

ABSTRACT. We present a moduli space for all hyperbolic basic sets of diffeomorphisms on surfaces that have an invariant measure that is absolutely continuous with respect to Hausdorff measure. To do this we introduce two new invariants: the measure solenoid function and the cocycle-gap pair. We extend the eigenvalue formula of A. N. Livšic and Ja. G. Sinai for Anosov diffeomorphisms which preserve an absolutely continuous measure to hyperbolic basic sets on surfaces which possess an invariant measure absolutely continuous with respect to Hausdorff measure. We characterise the Lipschitz conjugacy classes of such hyperbolic systems in a number of ways, for example, in terms of eigenvalues of periodic points and Gibbs measures.

CONTENTS

1. Introduction	2
2. Hyperbolic diffeomorphisms	9
3. Solenoid functions	14
4. Self-renormalisable structures	20
5. Measure solenoid functions	24
6. Natural geometric measures	29
7. Cocycle-gap pairs	37
8. Realisations of Gibbs measures	44
9. Eigenvalues	49
10. Invariant Hausdorff measures	55
References	58

1. INTRODUCTION

We say that (f, Λ) is a C^{1+} *hyperbolic diffeomorphism* if it has the following properties:

- (i) $f : M \rightarrow M$ is a $C^{1+\alpha}$ diffeomorphism of a compact surface M with respect to a $C^{1+\alpha}$ structure $\mathcal{C}_f$ on M , for some $\alpha > 0$;
- (ii) Λ is a hyperbolic invariant subset of M ; and
- (iii) $f|_\Lambda$ is topologically transitive and that Λ has a local product structure.

We denote by $\mathcal{T}(f, \Lambda)$ the set of all C^{1+} hyperbolic diffeomorphisms (g, Λ_g) such that (g, Λ_g) and (f, Λ) are topologically conjugated by a homeomorphism $h_{f,g}$ (see Section 2.5). We allow both the case where $\Lambda = M$ and the case where Λ is a proper subset of M . If $\Lambda = M$ then f is Anosov and M is a torus [9, 19]. The best known examples where Λ is a proper subset of M are the Smale horseshoes, and the codimension one attractors such as the Plykin attractor and the derived-Anosov diffeomorphisms.

For every $g \in \mathcal{T}(f, \Lambda)$, we denote by $\delta_{g,s}$ (resp. $\delta_{g,u}$) the Hausdorff dimension of the local stable (resp. local unstable) leaves of g intersected with Λ . Let $\lambda_{g,s}(x)$ and $\lambda_{g,u}(x)$ denote the stable and unstable eigenvalues of the periodic orbit of g containing a point x . A. N. Livšic and Ja. G. Sinai [13] proved that an Anosov diffeomorphism g has an invariant measure that is absolutely continuous with respect to Lebesgue measure if, and only if, $\lambda_{g,s}(x)\lambda_{g,u}(x) = 1$ for every periodic point x . We extend the theorem of A. N. Livšic and Ja. G. Sinai to C^{1+} hyperbolic diffeomorphisms with hyperbolic sets on surfaces such as Smale horseshoes and codimension one attractors.

Theorem 1.1. *A C^{1+} hyperbolic diffeomorphism $g \in \mathcal{T}(f, \Lambda)$ has a g -invariant probability measure which is absolutely continuous to the Hausdorff measure on Λ_g if and only if for every periodic point x of $g|_{\Lambda_g}$,*

$$\lambda_{g,s}(x)^{\delta_{g,s}} \lambda_{g,u}(x)^{\delta_{g,u}} = 1.$$

The proof of all theorems stated in the introduction are given in Section 10.

Since (f, Λ) is a C^{1+} hyperbolic diffeomorphism it admits a Markov partition $\mathcal{R} = \{R_1, \dots, R_k\}$. This implies the existence of a two-sided subshift $\tau : \Theta \rightarrow \Theta$ of finite type Θ in the symbol space $\{1, \dots, k\}^{\mathbb{Z}}$, and an inclusion $i : \Theta \rightarrow \Lambda$ such that (a) $f \circ i = i \circ \tau$ and (b) $i(\Theta_j) = R_j$ for every $j = 1, \dots, k$, where Θ_j is the cylinder containing all words $\dots \varepsilon_{-1} \varepsilon_0 \varepsilon_1 \dots \in \Theta$ with $\varepsilon_0 = j$. For every $g \in \mathcal{T}(f, \Lambda)$, the inclusion $i_g = h_{f,g} \circ i : \Theta \rightarrow \Lambda_g$ is such that $g \circ i_g = i_g \circ \tau$. We call such a map $i_g : \Theta \rightarrow \Lambda_g$ a *marking* of (g, Λ_g) .

Definition 1.1. *If $g \in \mathcal{T}(f, \Lambda)$ is a C^{1+} hyperbolic diffeomorphism as above and ν is a Gibbs measure on Θ then we say that (g, Λ_g, ν) is a Hausdorff realisation of ν if $(i_g)_* \nu$ is absolutely continuous with respect to the Hausdorff measure on Λ_g . If this is the case then we will often just say that ν is a Hausdorff realisation for (g, Λ_g) .*

We note that if $g \in \mathcal{T}(f, \Lambda)$ the Hausdorff measure on Λ_g exists and is unique. However, a Hausdorff realisation need not exist for (g, Λ_g) .

Let $\mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ be the set of all C^{1+} hyperbolic diffeomorphisms (g, Λ_g) in $\mathcal{T}(f, \Lambda)$ such that (i) $\delta_{g,s} = \delta_s$ and $\delta_{g,u} = \delta_u$; (ii) there is a g -invariant measure μ_g on Λ_g which is absolutely continuous with respect to the Hausdorff measure on Λ_g . We denote by $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ the subset of all C^{1+} -realisations of a Gibbs measure ν in $\mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$.

De la Llave, Marco and Moriyon [14, 15, 17, 18] have shown that the set of stable and unstable eigenvalues of all periodic points is a complete invariant of the C^{1+} conjugacy classes of Anosov diffeomorphisms. We extend their result to the sets $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$.

Theorem 1.2. *(i) Any two elements of $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ have the same set of stable and unstable eigenvalues and these sets are a complete invariant of $[\nu]$ in the sense that if $g_1, g_2 \in \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ have the same eigenvalues if, and only if, they are in the same subset $[\nu]$.*

(ii) The map $\nu \rightarrow [\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ gives a 1–1 correspondence between C^{1+} -Hausdorff realisable Gibbs measures ν and Lipschitz conjugacy classes in $\mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$.

In Theorem 9.6, we also prove that the set of stable and unstable eigenvalues of all periodic orbits of a C^{1+} hyperbolic diffeomorphism $g \in \mathcal{T}(f, \Lambda)$ is a complete invariant of each Lipschitz conjugacy class. We note that for Anosov diffeomorphisms every Lipschitz conjugacy class is a C^{1+} conjugacy class. This can be proved by combining Remark 8.1 with Lemmas 4.2 and 8.1.

Remark 1.3. *We have restricted our discussion to Gibbs measures because it follows from Theorem 1.2 that, if $g \in \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ has a g -invariant measure μ which is absolutely continuous with respect to the Hausdorff measure then μ is a C^{1+} -Hausdorff realisation of a Gibbs measure ν so that $\mu = (i_g)_*\nu$.*

E. Cawley [3] characterised all C^{1+} -Hausdorff realisable Gibbs measures as Anosov diffeomorphisms using cohomology classes on the torus. While it is possible that her cocycles could give enough information to characterise other hyperbolic systems on surfaces up to lippeomorphism, it is clear that they cannot encode enough for C^{1+} conjugacy because, for example, they do not encode enough information about gaps and so do not determine the smooth structure of stable leaves in the case where they are Cantor sets. To deal with all these cases in an integrated way we use measure solenoid functions and gap-cocycle pairs to classify C^{1+} -Hausdorff realisable Gibbs measures of all C^{1+} hyperbolic diffeomorphisms on surfaces.

The *stable and unstable measure solenoid functions* are built from the Gibbs measure as we show in Section 5.4. For Anosov diffeomorphisms, the domains msol^s and msol^u of the stable and unstable measure solenoid functions are dense subsets of finite disjoint unions of closed intervals. Define a stable leaf segment of a Markov rectangle to be a segment of a stable leaf crossing the Markov rectangle (see Section 2.3, Section 2.6, Section 2.7 and Figure 2). Every point in msol^s consists of a pair (I, J) of adjacent stable spanning leaf segments of Markov rectangles which are not contained in a stable global leaf containing a stable boundary of a Markov rectangle. The stable measure solenoid function $\sigma_{\nu,s}$ associates to each pair (I, J) the ratio between the measure of J and the measure of I computed with respect to the conditional measure, determined by the Gibbs measure ν , of a stable leaf containing I and J . The construction of the unstable solenoid function $\sigma_{\nu,u}$ is similar. In Section 3.7, we define a *boundary condition* which consists of a finite set of simple algebraic equalities that the continuous extensions of the stable and unstable measure solenoid functions have to satisfy, for the corresponding Gibbs measures to be C^{1+} -Hausdorff realisable in the Anosov case. This is necessary because in this case the Markov rectangles have common boundaries along the stable and unstable foliations. We note that in the Anosov case, the Lebesgue measure is the Hausdorff measure.

Theorem 1.4. *(Anosov diffeomorphisms) Suppose that f is a C^{1+} Anosov diffeomorphism of the torus Λ . Fix a Gibbs measure ν on Θ . Then the following statements are equivalent:*

- (i) *The set $\nu, [\nu] \subset \mathcal{T}_{f,\Lambda}(1, 1)$ is non-empty and is precisely the set of $g \in \mathcal{T}_{f,\Lambda}(1, 1)$ such that (g, Λ_g, ν) is a C^{1+} Hausdorff realisation. In this case $\mu = (i_g)_*\nu$ is absolutely continuous with respect to Lebesgue measure.*
- (ii) *The stable measure solenoid function $\sigma_{\nu,s} : \text{msol}^s \rightarrow \mathbb{R}^+$ has a non-vanishing Hölder continuous extension to the closure of msol^s satisfying the boundary condition.*
- (iii) *The unstable measure solenoid function $\sigma_{\nu,u} : \text{msol}^u \rightarrow \mathbb{R}^+$ has a non-vanishing Hölder continuous extension to the closure of msol^s satisfying the boundary condition.*

The treatment of codimension one attractors has a number of extra-dificulties due to the fact that the invariant set Λ is locally a Cartesian product of a Cantor set with an interval but the stable

and unstable measure solenoid functions are built in a similar way to the construction for Anosov diffeomorphisms. In the case of codimension one attractors, the continuous extension of the stable measure solenoid functions have to satisfy the *cylinder-cylinder condition* for the corresponding Gibbs measures to be C^{1+} -Hausdorff realisable (see Section 5.4). The cylinder-cylinder condition, like the boundary condition, consists of a finite set of simple algebraic equalities and is needed because the Markov rectangles have common boundaries along the stable laminations. Hence, the cylinder-cylinder condition just applies to the stable measure function.

Theorem 1.5. (*Codimension one attractors*) Suppose that f is a C^{1+} surface diffeomorphism and Λ is a codimension one hyperbolic attractor. Fix a Gibbs measure ν on Θ . Then the following statements are equivalent:

- (i) For all $0 < \delta_s < 1$, $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, 1)$ is non-empty and is precisely the set of $g \in \mathcal{T}_{f,\Lambda}(\delta_s, 1)$ such that (g, Λ_g, ν) is a C^{1+} Hausdorff realisation. In this case $\mu = (i_g)_*\nu$ is absolutely continuous with respect to the Hausdorff measure on Λ_g .
- (ii) The stable measure solenoid function $\sigma_{\nu,s} : \text{msol}^s \rightarrow \mathbb{R}^+$ has a non-vanishing Hölder continuous extension to the closure of msol^s satisfying the cylinder-cylinder condition.
- (iii) The unstable measure solenoid function $\sigma_{\nu,u} : \text{msol}^u \rightarrow \mathbb{R}^+$ has a non-vanishing Hölder continuous extension to the closure of msol^u .

In the case of Smale horseshoes, there are no extra conditions that the measure solenoid functions have to satisfy for the corresponding Gibbs measures to be C^{1+} -Hausdorff realisable.

Theorem 1.6. (*Smale horseshoes*) Suppose that (f, Λ) is a Smale horseshoe and ν is a Gibbs measure on Θ . Then for all $0 < \delta_s, \delta_u < 1$, $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ is non-empty and is precisely the set of $g \in \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ such that (g, Λ_g, ν) is a C^{1+} Hausdorff realisation. In this case $\mu = (i_g)_*\nu$ is absolutely continuous with respect to the Hausdorff measure on Λ_g .

Using Theorems 1.2-1.5, for $\iota \in \{s, u\}$, we prove that the map $\nu \rightarrow \sigma_{\nu,\iota}$ gives a one-to-one correspondence between the subsets $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ and the measure solenoid functions $\sigma_{g,\iota}$ satisfying the conditions indicated in the above Theorems 1.4 and 1.5. By Theorem 1.2, we have that the sets $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ are precisely the Lipschitz conjugacy classes contained in $\mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$.

Theorem 1.7. *The measure solenoid functions determine a pair of infinite-dimensional metric spaces, that we denote by SOL^s and SOL^u , which parametrize all Lipschitz conjugacy classes $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$.*

The scaling functions, presented by M. Feigenbaum in [7, 8] and D. Sullivan in [33], and the solenoid functions, presented in [21, 24, 28] (see also Section 1.3 and Section 3), are used to classify all C^{1+} conjugacy classes of expanding maps on train-tracks and of hyperbolic diffeomorphisms for a given a topological conjugacy class. Both scaling function and solenoid function are complete invariants of the smooth structure but the solenoid functions have the great advantage that, unlike the scaling functions, one knows which solenoid functions occur as the solenoid functions of C^{1+} expanding maps.

In Section 7, we introduce the *stable and unstable cocycle-gap pairs* (γ_s, J_s) and (γ_u, J_u) which allow us to parametrize the C^{1+} conjugacy classes inside of each Lipschitz conjugacy class $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$. The cocycle-gap pair (γ, J) consists of a gap function γ and a measure-length ratio cocycle function J (see Definitions 7.1 and 7.2). The domain of a stable gap function is the set of all pairs (ξ_1, ξ_2) with the following properties: (a) the stable leaf segments ξ_1 and ξ_2 intersect the invariant set Λ just in their end points; (b) there is a stable leaf K of a Markov rectangle which contains $f\xi_1$ and $f\xi_2$. The *stable gap function* γ is a Hölder continuous function satisfying the following algebraic equality $\gamma(\xi_1 : \xi_2) = \gamma(\xi_1 : \xi_3)\gamma(\xi_3 : \xi_2)$. (We use the notation $\xi_1 : \xi_2$

rather than ξ_1, ξ_2 just to emphasise that γ will be measuring ratios.) The domain of a stable measure-length ratio cocycle is the set of all stable spanning leaves of Markov rectangles. A *stable measure-length ratio cocycle* is a function $J = \kappa/(\kappa \circ f)$ where κ is a positive Hölder continuous function satisfying a set of algebraic inequalities given by (7.1). The unstable gap functions and measure-length ratio cocycle are defined similarly. In the case of Smale horseshoes the domains of the gap functions and of the measure-length ratio cocycle functions are Cantor sets. In the case of codimension one attractors the domains of the stable gap functions and of the stable measure-length ratio cocycle functions are a finite union of intervals, and the stable cocycle-gap pairs have to satisfy the *cocycle-gap property* (see Definition 7.4 and Lemma 7.9) which is due to the fact that the Markov rectangles have common boundaries along the stable laminations. The stable and unstable cocycle-gap pairs give rise to an infinite dimensional metric space, that we denote respectively by $\mathcal{JG}^s(\nu, \delta_s)$ and $\mathcal{JG}^u(\nu, \delta_u)$.

Theorem 1.8. (i) (Smale horseshoes) *There is a natural map*

$$g \rightarrow (\gamma_s(g), J_s(g), \gamma_u(g), J_u(g))$$

which gives a one-to-one correspondence between C^{1+} conjugacy classes of diffeomorphisms g contained in $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ and stable and unstable cocycle-gap pairs contained in $\mathcal{JG}^s(\nu, \delta_s) \times \mathcal{JG}^u(\nu, \delta_u)$.

(ii) (Codimension one attractors) *There is a natural map $g \rightarrow (\gamma_s(g), J_s(g))$ which gives a one-to-one correspondence between C^{1+} conjugacy classes of diffeomorphisms g contained in $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, 1)$ and stable cocycle-gap pairs contained in $\mathcal{JG}^s(\nu, \delta_s)$.*

1.1. Self-renormalisable structures. In Section 4, we construct C^{1+} *stable and unstable self-renormalisable structures on train-tracks*. The train-tracks are a form of optimal local leaf-quotient space of the stable and unstable laminations of Λ . Locally, these train-tracks are just the quotient space of stable or unstable leaves within a Markov rectangle, but globally the identification of leaves common to two more than one rectangle gives a non-trivial structure and introduces junctions. They are characterised by being the compact quotient on which the Markov map induced by the action of f is continuous with the minimal number of identifications. A smooth structure on the stable or unstable leaves of Λ induces a smooth structure on the corresponding train-tracks and *vice-versa*. Now, we use that the holonomies of codimension one hyperbolic systems are C^{1+} (see [25]), and so the holonomies also project in the train-tracks and together with the Markov maps give rise to what we call self-renormalisable C^{1+} structures (see Figures 1, 8 and 9).

Remark 1.9. *These structures are called self-renormalisable because the train track $\mathbf{B}$ has defined on it both a Markov map m and a pseudo group of holonomy maps $\{h\}$. For the train tracks arising from Anosov diffeomorphisms of the torus $\{h\}$ can be identified with a C^{1+} diffeomorphism g of the circle and the Markov map m defines a renormalisation of g which agrees with the usual one ([29, 30]). The renormalised map Rg is C^{1+} conjugate to g . Thus these C^{1+} self-renormalisable structures are in one-to-one correspondance with C^{1+} fixed points of the the circle map renormalisation operator. It then follows from the results of this paper that these fixed points are in one-to-one correspondance with those Anosov diffeomorphisms of the the torus which preserve an a measure that is absolutely continuous with respect to Lebesgue measure. For train tracks arising from non-Anosov diffeomorphisms one gets other interesting renormalisation structures, for example, for interval exchange maps.*

In Section 4.7, we prove the following useful equivalence between 2-dimensional dynamics and 1-dimensional dynamics.

Theorem 1.10. *There is a natural map $g \rightarrow (\mathcal{S}_s(g), \mathcal{S}_u(g))$ which gives a one-to-one correspondence between C^{1+} conjugacy classes in $\mathcal{T}(f, \Lambda)$ and pairs of stable and unstable C^{1+} self-renormalisable structures.*




FIGURE 1. This figure illustrates a (unstable) train track for the Anosov map $g : \mathbb{R}^2 \setminus (\mathbb{Z}v \times \mathbb{Z}w) \rightarrow \mathbb{R}^2 \setminus (\mathbb{Z}v \times \mathbb{Z}w)$ defined by $g(x, y) = (x + y, y)$. The rectangles A and B are the Markov rectangles and the vertical arrows show paths along unstable manifolds from A to A and from B to A . The train track is represented by the pair of circles and the curves below it show the smooth paths through the junction of the two circles which arise from the smooth paths between the rectangles A and B along unstable manifolds. Note that there is no smooth path from B to B even though in this representation of the train track it looks as though there ought to be. This is because there is no unstable manifold running directly from the rectangle B to itself.

Hence, for a pair $(\mathcal{S}_s, \mathcal{S}_u)$ of C^{1+} self-renormalisable structures to be realisable by a C^{1+} hyperbolic diffeomorphism in $\mathcal{T}(f, \Lambda)$, the unstable C^{1+} self-renormalisable structure does not impose any restriction in the stable C^{1+} self-renormalisable structure, and *vice-versa*. The same is no longer true if we ask $g \in \mathcal{T}(f, \Lambda)$ to be a C^{1+} -Hausdorff realisation of a Gibbs measure as we describe in the next section.

1.2. Realisation of Gibbs measures on train-tracks. We are going to study the C^{1+} -Hausdorff realisations of Gibbs measures as self-renormalisable structures. Then we use this information to study the C^{1+} -Hausdorff realisations of Gibbs measures as C^{1+} hyperbolic diffeomorphisms, going like this from one-dimensional dynamics to two-dimensional dynamics.

Let $\pi_u : \Theta \rightarrow \Theta^u$ and $\pi_s : \Theta \rightarrow \Theta^s$ be respectively the natural projections on the left and right infinite words. Let $\tau_u : \Theta^u \rightarrow \Theta^u$ and $\tau_s : \Theta^s \rightarrow \Theta^s$ be the corresponding right and left shifts. Let $\iota \in \{s, u\}$ where s denotes stable and u denotes unstable. Let us denote by $\mathbf{B}^\iota$ the ι train-track and by $i_\iota : \Theta^\iota \rightarrow \mathbf{B}^\iota$ the natural marking induced by $i : \Theta \rightarrow \Lambda$ (see Section 4.5). A τ -invariant measure ν on Θ determines a unique τ_ι -invariant measure $\nu_\iota = (\pi_\iota)_* \nu$ on Θ^ι . Conversely, a τ_ι -invariant measure ν_ι on Θ^ι has a unique natural extension to a τ -invariant measure ν on Θ . We say that a τ_ι -invariant measure ν_ι is a *Gibbs measure* if its natural extension ν is a Gibbs measure on Θ . A C^{1+} ι self-renormalisable structure $\mathcal{S}_\iota$ is a C^{1+} -Hausdorff realisation of a Gibbs measure ν on Θ if, for every chart (e, U) of the self-renormalisable structure $\mathcal{S}_\iota$, the pushforward $(e \circ i_\iota)_* \nu_\iota$ of the measure ν_ι is a measure absolutely continuous with respect to the Hausdorff measure of the set $e(U)$. Let us denote by $\delta(\mathcal{S}_\iota)$ the Hausdorff dimension of the set $e(U)$ for every chart (e, U) of the smooth structure of $\mathcal{S}_\iota$. Clearly, this is independent of the chart (e, U) . We denote by $\mathcal{D}^\iota(\nu, \delta)$ the set of all C^{1+} ι self-renormalisable structures $\mathcal{S}_\iota$ with $\delta(\mathcal{S}_\iota) = \delta$ and which are C^{1+} -Hausdorff realisations

of the Gibbs measure ν on Θ . By Theorem 6.1, every C^{1+} ι self-renormalisable structure $\mathcal{S}_\iota$ is a C^{1+} -Hausdorff realisation of an unique Gibbs measure $\nu_{\mathcal{S}_\iota}$ on Θ .

Theorem 1.11. *The map $g \rightarrow (\mathcal{S}_s(g), \mathcal{S}_u(g))$ gives a 1-1 correspondence between C^{1+} conjugacy classes in $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ and pairs in $\mathcal{D}^s(\nu, \delta_s) \times \mathcal{D}^u(\nu, \delta_u)$.*

Hence, if $g \in \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ then $\delta(\mathcal{S}_s(g)) = \delta_s$ and $\delta(\mathcal{S}_u(g)) = \delta_u$. Let $\mathcal{S}_\iota$ be a C^{1+} ι self-renormalisable structure. If $\delta(\mathcal{S}_\iota) = 1$ we call $\mathbf{B}^\iota$ a *no-gap train-track*. If $0 < \delta(\mathcal{S}_\iota) < 1$ we call $\mathbf{B}^\iota$ a *gap train-track*. Let ι' denote the element of $\{s, u\}$ which is not $\iota \in \{s, u\}$.

Theorem 1.12. *Let $\mathbf{B}^s$ and $\mathbf{B}^u$ be the stable and unstable train-tracks determined by a C^{1+} hyperbolic diffeomorphism (f, Λ) . The set $\mathcal{D}^\iota(\nu, \delta_\iota)$ is non-empty if, and only if, the ι -measure solenoid function $\sigma_\nu : \text{msol}^\iota \rightarrow \mathbb{R}^+$ of the Gibbs measure ν has the following properties:*

- (i) *If $\mathbf{B}^\iota$ and $\mathbf{B}^{\iota'}$ are no-gap train-tracks then σ_ν has a non-vanishing Hölder continuous extension to the closure of msol^ι satisfying the boundary condition.*
- (ii) *If $\mathbf{B}^\iota$ is a no-gap train-track and $\mathbf{B}^{\iota'}$ is a gap train-track then σ_ν has a non-vanishing Hölder continuous extension to the closure of msol^ι .*
- (iii) *If $\mathbf{B}^\iota$ is a gap train-track and $\mathbf{B}^{\iota'}$ is a no-gap train-track then σ_ν has a non-vanishing Hölder continuous extension to the closure of msol^ι satisfying the cylinder-cylinder condition.*
- (iv) *If $\mathbf{B}^\iota$ and $\mathbf{B}^{\iota'}$ are gap train-tracks then σ_ν does not have to satisfy any extra-condition.*

Furthermore, $\mathcal{D}^\iota(\nu, \delta_\iota) \neq \emptyset$ if, and only if, $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'}) \neq \emptyset$

By Theorem 10.1, the set of all ι -measure solenoid functions σ_ν with the properties indicated in Theorem 1.12 determine an infinite dimensional metric space SOL^ι which gives a nice parametrization of all Lipschitz conjugacy classes $\mathcal{D}^\iota(\nu, \delta)$ of C^{1+} self-renormalisable structures $\mathcal{S}_\iota$ with a given Hausdorff dimension $\delta(\mathcal{S}_\iota) = \delta$. We have used the same notation SOL^ι as in Section 1 above because we will show that they are effectively the same sets.

Theorem 1.13. *Let us suppose that $\mathcal{D}^\iota(\nu, \delta) \neq \emptyset$.*

- (i) *(Flexibility) If $\mathbf{B}^\iota$ is a gap train-track then $\mathcal{D}^\iota(\nu, \delta)$ is an infinite dimensional space parametrized by cocycle-gap pairs contained in $\mathcal{JG}^\iota(\nu, \delta)$.*
- (ii) *(Rigidity) If $\mathbf{B}^\iota$ is a no-gap train-track then $\mathcal{D}^\iota(\nu, 1)$ consists of a single C^{1+} self-renormalisable structure.*

By Lemma 9.1, each set $\mathcal{D}^\iota(\nu, \delta)$ is either empty or a Lipschitz conjugacy class. Hence, if $\mathbf{B}^\iota$ is a no-gap train-track then the Lipschitz conjugacy class consists of a single C^{1+} self-renormalisable structure. Furthermore, by Lemma 9.5, the set of eigenvalues of all periodic orbits of $\mathcal{S}_\iota$ is a complete invariant of each set $\mathcal{D}^\iota(\nu, \delta)$ (see also [34]).

Theorem 1.14. *(Rigidity) If $\delta_\iota = 1$, the mapping $g \rightarrow \mathcal{S}_{\iota'}(g)$ gives a 1-1 correspondence between C^{1+} conjugacy classes in $[\nu] \subset \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ and C^{1+} self-renormalisable structures in $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'})$.*

Hence, the map $g \rightarrow \mathcal{S}_{\iota'}(g)$ gives rise to the following one-to-one correspondences. For Anosov diffeomorphisms (f, Λ) , there is an one-to-one correspondence between (i) C^{1+} conjugacy classes of hyperbolic diffeomorphisms $g \in \mathcal{T}(f, \Lambda)$ which are C^{1+} -Hausdorff realisations of a Gibbs measure; (ii) C^{1+} unstable self-renormalisable structures; and (iii) C^{1+} stable self-renormalisable structures. For codimension one attractors (f, Λ) , there is a one-to-one correspondence between (i) C^{1+} conjugacy classes of hyperbolic diffeomorphisms $g \in \mathcal{T}(f, \Lambda)$ which are C^{1+} -Hausdorff realisations of a Gibbs measure; and (ii) C^{1+} stable self-renormalisable structures.

1.3. Solenoid functions and Gibbs measures. In Section 3, we introduce *HR-structures* (HR for Hölder-ratios). These associate an affine structure to each stable and unstable leaf segment in such a way that these vary Hölder continuously with the leaf and are kept invariant by the dynamics of f . As we will describe the HR-structures are in one-to-one correspondence with the C^{1+} conjugacy classes of C^{1+} hyperbolic diffeomorphisms in $\mathcal{T}(f, \Lambda)$. The HR-structures are characterized by a pair (r_s, r_u) of ratio functions. The ratio functions r_s and r_u are independent one from another. However, if we ask that the C^{1+} hyperbolic diffeomorphism have an associated geometric measure then we will see later that the ratio function r_u imposes restrictions on the ratio function r_s and vice-versa.

For $\iota \in \{s, u\}$, we construct a topological set $\mathbf{S}^\iota$ which is either isomorphic to a finite union of closed intervals on the real line or to an embedded Cantor set on the real line. The ratio functions r_ι when restricted to the set $\mathbf{S}^\iota$ are Hölder continuous functions $\sigma_\iota = r_\iota|_{\mathbf{S}^\iota}$ completely characterised by a finite set of properties that we explain also in Section 3. We call the functions σ_ι *solenoid functions*. They form an infinite dimensional metric space which parametrizes the C^{1+} conjugacy classes of C^{1+} hyperbolic diffeomorphisms contained in $\mathcal{T}(f, \Lambda)$ (see Corollary 3.6).

We say that D is a *stable leaf n -cylinder segment* of Λ if $f^{-n}D$ is a stable spanning leaf segment of a Markov rectangle. Let E_D be a stable spanning leaf segment of a Markov rectangle such that $D \subset E_D$. Let ν be a Gibbs measure and $\mu = i_*\nu$ the corresponding invariant measure in Λ . Let $\rho(D \cap \Lambda, E_D \cap \Lambda)$ be the ratio between the measure of $D \cap \Lambda$ and the measure of $E_D \cap \Lambda$ with respect to the conditional measure of μ in E_D . The δ -*stable bounded solenoid class* of a Gibbs measure ν is the set of all stable solenoid functions σ with the following property: There is $C = C(\sigma) > 0$ such that for every stable leaf n -cylinder segment of Λ we have

$$|\delta_\iota \log r(D \cap \Lambda, E_D \cap \Lambda) - \log \rho(D \cap \Lambda, E_D \cap \Lambda)| < C,$$

where r is the stable ratio function determined by the stable solenoid function σ (see also Definition 8.2). The δ -unstable bounded solenoid class of a Gibbs measure ν is defined in a similar way.

- Theorem 1.15.**
- (i) *There is a natural map $g \rightarrow (\sigma_s(g), \sigma_u(g))$ which gives a one-to-one correspondence between C^{1+} conjugacy classes of C^{1+} hyperbolic diffeomorphisms $g \in \mathcal{T}(\nu, \delta_s, \delta_u)$ and pairs $(\sigma_s(g), \sigma_u(g))$ of stable and unstable solenoid functions such that, for ι equal to s and u , $\sigma_\iota(g)$ is contained in the δ_ι -bounded solenoid equivalence class of ν .*
 - (ii) *There is a natural map $\mathcal{S}_\iota \rightarrow \sigma_{\mathcal{S}_\iota}$ which gives a one-to-one correspondence between C^{1+} self-renormalisable structures $\mathcal{S}_\iota$ contained in $\mathcal{D}^\iota(\nu, \delta_\iota)$ and ι -solenoid functions $\sigma_{\mathcal{S}_\iota}$ contained in the δ_ι -bounded equivalence class of ν .*
 - (iii) *Let us suppose that $\mathcal{D}^\iota(\nu, \delta_\iota) \neq \emptyset$.*
 - (a) *(Rigidity) If $\delta_\iota = 1$ then the δ_ι -bounded solenoid equivalence class of ν is a singleton consisting in the continuous extension of the ι measure solenoid function $\sigma_{\nu, \iota}$ to $\mathbf{S}^\iota$.*
 - (b) *(Flexibility) If $0 < \delta_\iota < 1$ then the δ_ι -bounded solenoid equivalence class of ν is an infinite dimensional space of solenoid functions.*

In Section 7, we use the cocycle-gap pairs to construct explicitly the solenoid functions in the δ_ι -bounded equivalence classes of the C^{1+} -Hausdorff realisable Gibbs measures ν .

By Lemma 4.1, given an ι -solenoid function σ_ι there is a unique C^{1+} self-renormalisable structure $\mathcal{S}_\iota$ such that $\sigma_\iota = \sigma_{\mathcal{S}_\iota}$ and, by Theorem 6.1 and Lemma 6.3, there is a unique C^{1+} -Hausdorff realisable Gibbs measure $\nu = \nu_{\sigma_\iota}$ such that $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota)$ with $\delta_\iota = \delta(\mathcal{S}_\iota)$. Hence, by Theorem 1.15 (ii), given an ι -solenoid function σ_ι there is a unique C^{1+} -Hausdorff realisable Gibbs measure ν such that σ_ι belongs to the δ_ι -bounded solenoid equivalence class of ν .

Theorem 1.16. *Given an ι -solenoid function σ_ι and $0 < \delta_{\iota'} \leq 1$, there is a unique Gibbs measure ν and a unique $\delta_{\iota'}$ -bounded equivalence class of ν consisting of ι' -solenoid functions $\sigma_{\iota'}$ such that the C^{1+} conjugacy class of hyperbolic diffeomorphisms $g \in \mathcal{T}_{f,\Lambda}(\delta_s, \delta_u)$ determined by the pair (σ_s, σ_u) have an invariant measure $\mu = (i_g)_*\nu$ absolutely continuous with respect to the Hausdorff measure.*

Putting together Theorem 1.15 and Theorem 1.16, we obtain the following implications:

- (i) (Flexibility for Smale horseshoes) For $\iota = s$ and u , given a ι -solenoid function σ_ι there is an infinite dimensional space of solenoid functions $\sigma_{\iota'}$ such that the C^{1+} hyperbolic Smale horseshoes determined by the pairs (σ_s, σ_u) have an invariant measure μ absolutely continuous with respect to the Hausdorff measure.
- (ii) (Rigidity for Anosov diffeomorphisms) For $\iota = s$ and u , given an ι -solenoid function σ_ι there is a unique ι' -solenoid function such that the C^{1+} Anosov diffeomorphisms determined by the pair (σ_s, σ_u) has an invariant measure μ absolutely continuous with respect to Lebesgue.
- (iii) (Flexibility for codimension one attractors) Given an unstable solenoid function σ_u there is an infinite dimensional space of stable solenoid functions σ_s such that the C^{1+} hyperbolic codimension one attractors determined by the pairs (σ_s, σ_u) have an invariant measure μ absolutely continuous with respect to the Hausdorff measure.
- (iv) (Rigidity for codimension one attractors) Given an s -solenoid function σ_s there is a unique unstable solenoid function σ_u such that the C^{1+} hyperbolic codimension one attractors determined by the pair (σ_s, σ_u) have an invariant measure μ absolutely continuous with respect to the Hausdorff measure using non-zero stable and unstable pressures.

In this paper we prove, in fact, a generalized version of the results presented in the introduction. The theory that we develop here studies the properties and classifies all the Gibbs measures which are C^{1+} -realisable as natural geometric measures of C^{1+} self-renormalisable structures (see Definition 6.1) and of C^{1+} hyperbolic diffeomorphisms (see Definition 8.3). The set of natural geometric measures contains the invariant measures which are absolutely continuous with respect to Hausdorff measure.

The proof of all theorems stated in the introduction are given in Section 10.

2. HYPERBOLIC DIFFEOMORPHISMS

In this section, we present some basic facts on hyperbolic dynamics, that we include for clarity of the exposition.

2.1. Interval notation. We also use the notation of interval arithmetic for some inequalities where:

- (i) if I and J are intervals then $I + J$, $I.J$ and I/J have the obvious meaning as intervals,
- (ii) if $I = \{x\}$ then we often denote I by x , and
- (iii) $I \pm \varepsilon$ denotes the interval consisting of those x such that $|x - y| < \varepsilon$ for all $y \in I$.

By $\phi(n) \in 1 \pm \mathcal{O}(\nu^n)$ we mean that there exists a constant $c > 0$ depending only upon explicitly mentioned quantities such that for all $n \geq 0$, $1 - c\nu^n < \phi(n) < 1 + c\nu^n$. By $\phi(n) = \mathcal{O}(\nu^n)$ we mean that there exists a constant $c \geq 1$ depending only upon explicitly mentioned quantities such that for all $n \geq 0$, $c^{-1}\nu^n \leq \phi(n) \leq c\nu^n$.

2.2. Stable and unstable superscripts. Throughout the paper we will use the following notation: we use ι to denote an element of the set $\{s, u\}$ of the stable and unstable superscripts and ι' to denote the element of $\{s, u\}$ that is not ι . In the main discussion we will often refer to objects

which are qualified by ι such as, for example, an ι -leaf: This means a leaf which is a leaf of the stable lamination if $\iota = s$, or a leaf of the unstable lamination if $\iota = u$. In general the meaning should be quite clear.

We define the map $f_\iota = f$ if $\iota = u$ or $f_\iota = f^{-1}$ if $\iota = s$.

2.3. Leaf segments. Let d be a metric on M . For $\iota \in \{s, u\}$, if $x \in \Lambda$ we denote the local ι -manifolds through x by

$$W^\iota(x, \varepsilon) = \{y \in M : d(f_\iota^{-n}(x), f_\iota^{-n}(y)) \leq \varepsilon, \text{ for all } n \geq 0\}.$$

By the Stable Manifold Theorem (see [11]), these sets are respectively contained in the stable and unstable immersed manifolds

$$W^\iota(x) = \bigcup_{n \geq 0} f_\iota^n(W^\iota(f_\iota^{-n}(x), \varepsilon_0))$$

which are the image of a $C^{1+\gamma}$ immersion $\kappa_{\iota,x} : \mathbb{R} \rightarrow M$. An *open (resp. closed) full ι -leaf segment* I is defined as a subset of $W^\iota(x)$ of the form $\kappa_{\iota,x}(I_1)$ where I_1 is a non-empty open (resp. closed) subinterval in $\mathbb{R}$. An *open (resp. closed) ι -leaf segment* is the intersection with Λ of an open (resp. closed) full ι -leaf segment such that the intersection contains at least two distinct points. If the intersection is exactly two points we call this closed ι -leaf segment an *ι -leaf gap*. A *full ι -leaf segment* is either an open or closed full ι -leaf segment. An *ι -leaf segment* is either an open or closed ι -leaf segment. The *endpoints* of a full ι -leaf segment are the points $\kappa_{\iota,x}(u)$ and $\kappa_{\iota,x}(v)$ where u and v are the endpoints of I_1 . The *endpoints* of an ι -leaf segment I are the points of the minimal closed full ι -leaf segment containing I . The *interior* of an ι -leaf segment I is the complement of its boundary. In particular, an ι -leaf segment I has empty interior if, and only if, it is an ι -leaf gap. A map $c : I \rightarrow \mathbb{R}$ is an *ι -leaf chart* of an ι -leaf segment I if has an extension $c_E : I_E \rightarrow \mathbb{R}$ to a full ι -leaf segment I_E with the following properties: $I \subset I_E$ and c_E is a homeomorphism onto its image.

2.4. Smoothness. In this paper, when we say that a map, atlas or structure is C^r we include the case C^{k+} where k is a positive integer. For maps f this means that f is $C^{k+\alpha}$ for some $0 < \alpha < 1$, i.e. C^k with α -Hölder continuous k th-order derivatives. For an atlas or structure by C^{k+} we mean that each pair of charts in the atlas or structure are $C^{k+\alpha}$ compatible for some $0 < \alpha < 1$ where the α might depend upon the charts. In the case of an atlas, we suppose that (i) one can choose α to be independent of the charts and (ii) the overlap maps have $C^{k+\alpha}$ norm bounded independent of the charts considered. This is immediately verified if the number of charts contained in the C^{k+} atlas is finite. Thus a C^{k+} atlas is $C^{k+\alpha}$, for some $0 < \alpha < 1$. This is not the case for C^{k+} structures.

2.5. Topological and smooth conjugacies. Let (f, Λ) be a C^{1+} hyperbolic diffeomorphism. Somewhat unusually we also desire to highlight the C^{1+} structure on M in which f is a diffeomorphism. By a C^{1+} structure on M we mean a maximal set of charts with open domains in M such that the union of their domains cover M and whenever U is an open subset contained in the domains of any two of these charts i and j then the overlap map $j \circ i^{-1} : i(U) \rightarrow j(U)$ is $C^{1+\alpha}$, where $\alpha > 0$ depends on i, j and U . We note that by compactness of M , given such a C^{1+} structure on M , there is an atlas consisting of a finite set of these charts which cover M and for which the overlap maps are $C^{1+\alpha}$ compatible and uniformly bounded in the $C^{1+\alpha}$ norm, where $\alpha > 0$ just depends upon the atlas. We denote by $\mathcal{C}_f$ the C^{1+} structure on M in which f is a diffeomorphism. Usually one is not concerned with this as, given two such structures, there is a homeomorphism of M sending one onto the other and thus, from this point of view, all such structures can be identified. For our discussion it will be important to maintain the identity of the different smooth structures on M .

We say that a map $h : \Lambda_f \rightarrow \Lambda_g$ is a *topological conjugacy* between two C^{1+} hyperbolic diffeomorphisms (f, Λ_f) and (g, Λ_g) if there is a homeomorphism $h : \Lambda_f \rightarrow \Lambda_g$ with the following properties:

- (i) $g \circ h(x) = h \circ f(x)$ for every $x \in \Lambda_f$.
- (ii) The pull-back of the ι -leaf segments of g by h are ι -leaf segments of f .

We say that a topological conjugacy $h : \Lambda_f \rightarrow \Lambda_g$ is a *Lipschitz conjugacy* if h has a bi-Lipschitz homeomorphic extension to an open neighbourhood of Λ_f in the surface M (with respect to the C^{1+} structures $\mathcal{C}_f$ and $\mathcal{C}_g$, respectively).

Similarly, we say that a topological conjugacy $h : \Lambda_f \rightarrow \Lambda_g$ is a $C^{1+\alpha}$ *conjugacy* if h has a $C^{1+\alpha}$ diffeomorphic extension to an open neighbourhood of Λ_f in the surface M , for some $\alpha > 0$.

Our approach is to fix a C^{1+} hyperbolic diffeomorphism (f, Λ) and consider C^{1+} hyperbolic diffeomorphism (g_1, Λ_{g_1}) topologically conjugate to (f, Λ) . The topological conjugacy $h : \Lambda \rightarrow \Lambda_{g_1}$ between f and g_1 extends to a homeomorphism H defined on a neighbourhood of Λ . Then, we obtain the new C^{1+} -realization (g_2, Λ_{g_2}) of f defined as follows: (i) the map $g_2 = H^{-1} \circ g_1 \circ H$; (ii) the basic set is $\Lambda_{g_2} = H^{-1}|\Lambda_{g_1}$; (iii) the C^{1+} structure $\mathcal{C}_{g_2}$ is given by the pull-back $(H)_* \mathcal{C}_{g_1}$ of the C^{1+} structure $\mathcal{C}_{g_1}$. From (i) and (ii), we get that $\Lambda_{g_2} = \Lambda$ and $g_2|_{\Lambda} = f$. From (iii), we get that g_2 is C^{1+} conjugated to g_1 . Hence, to study the conjugacy classes of C^{1+} hyperbolic diffeomorphisms (f, Λ) of f , we can just consider the C^{1+} hyperbolic diffeomorphisms (g, Λ_g) with $\Lambda_g = \Lambda$ and $g|_{\Lambda} = f|_{\Lambda}$, which we will do from now on for simplicity of our exposition.

2.6. Rectangles. Since Λ is a hyperbolic invariant set of a diffeomorphism $f : M \rightarrow M$, for $0 < \varepsilon < \varepsilon_0$ there is $\delta = \delta(\varepsilon) > 0$ such that, for all points $w, z \in \Lambda$ with $d(w, z) < \delta$, $W^u(w, \varepsilon)$ and $W^s(z, \varepsilon)$ intersect in a unique point that we denote by $[w, z]$. Since we assume that the hyperbolic set has a *local product structure*, we have that $[w, z] \in \Lambda$. Furthermore, the following properties are satisfied: (i) $[w, z]$ varies continuously with $w, z \in \Lambda$; (ii) the bracket map is continuous on a δ -uniform neighbourhood of the diagonal in $\Lambda \times \Lambda$; and (iii) whenever both sides are defined $f([w, z]) = [f(w), f(z)]$. Note that the bracket map does not really depend on δ provided it is sufficiently small.

Let us underline that it is a standing hypothesis that all the hyperbolic sets considered here have such a local product structure.

A *rectangle* R is a subset of Λ which is (i) closed under the bracket i.e. $x, y \in R \implies [x, y] \in R$, and (ii) proper i.e. is the closure of its interior in Λ . This definition imposes that a rectangle has always to be proper which is more restrictive than the usual one which only insists on the closure condition.

If ℓ^s and ℓ^u are respectively stable and unstable leaf segments intersecting in a single point then we denote by $[\ell^s, \ell^u]$ the set consisting of all points of the form $[w, z]$ with $w \in \ell^s$ and $z \in \ell^u$. We note that if the stable and unstable leaf segments ℓ and ℓ' are closed then the set $[\ell, \ell']$ is a rectangle. Conversely in this 2-dimensional situations, any rectangle R has a product structure in the following sense: for each $x \in R$ there are closed stable and unstable leaf segments of Λ , $\ell^s(x, R) \subset W^s(x)$ and $\ell^u(x, R) \subset W^u(x)$ such that $R = [\ell^s(x, R), \ell^u(x, R)]$. The leaf segments $\ell^s(x, R)$ and $\ell^u(x, R)$ are called *stable and unstable spanning leaf segments* for R (see Figure 2). For $\iota \in \{s, u\}$, we denote by $\partial \ell^\iota(x, R)$ the set consisting of the endpoints of $\ell^\iota(x, R)$, and we denote by $\text{int } \ell^\iota(x, R)$ the set $\ell^\iota(x, R) \setminus \partial \ell^\iota(x, R)$. The *interior* of R is given by $\text{int } R = [\text{int } \ell^s(x, R), \text{int } \ell^u(x, R)]$, and the *boundary* of R is given by $\partial R = [\partial \ell^s(x, R), \ell^u(x, R)] \cup [\ell^s(x, R), \partial \ell^u(x, R)]$.

2.7. Markov partitions. A *Markov partition* of f is a collection $\mathcal{R} = \{R_1, \dots, R_k\}$ of rectangles such that (i) $\Lambda \subset \bigcup_{i=1}^k R_i$; (ii) $R_i \cap R_j = \partial R_i \cap \partial R_j$ for all i and j ; (iii) if $x \in \text{int } R_i$ and $fx \in \text{int } R_j$ then

- (a) $f(\ell^s(x, R_i)) \subset \ell^s(fx, R_j)$ and $f^{-1}(\ell^u(fx, R_j)) \subset \ell^u(x, R_i)$




FIGURE 2. A rectangle.

(b) $f(\ell^u(x, R_i)) \cap R_j = \ell^u(fx, R_j)$ and $f^{-1}(\ell^s(fx, R_j)) \cap R_i = \ell^s(x, R_i)$.

The last condition means that $f(R_i)$ goes across R_j just once. In fact, it follows from condition (a) providing the rectangles R_j are chosen sufficiently small (see [16]). The rectangles making up the Markov partition are called *Markov rectangles*.

We note that there is a Markov partition $\mathcal{R}$ of f with the following *disjointness property* (see [2, 20, 32]):

- (i) if $0 < \delta_{f,s} < 1$ and $0 < \delta_{f,u} < 1$ then the stable and unstable leaf boundaries of any two Markov rectangles do not intersect.
- (ii) if $0 < \delta_{f,\iota} < 1$ and $\delta_{f,\iota'} = 1$ then the ι' -leaf boundaries of any two Markov rectangles do not intersect except, possibly, at their endpoints.

If $\delta_{f,s} = \delta_{f,u} = 1$, the disjointness property does not apply and so we consider that it is trivially satisfied for every Markov partition. For simplicity of our exposition, we will just consider Markov partitions satisfying the disjointness property.

2.8. Marking the invariant set Λ . The properties of the Markov partition $\mathcal{R} = \{R_1, \dots, R_k\}$ of f imply the existence of an unique two-sided subshift τ of finite type $\Theta = \Theta_A$ and a continuous surjection $i : \Theta \rightarrow \Lambda$ such that (a) $f \circ i = i \circ \tau$ and (b) $i(\Theta_j) = R_j$ for every $j = 1, \dots, k$. We call such a map $i : \Theta \rightarrow \Lambda$ a *marking of a C^{1+} hyperbolic diffeomorphism (f, Λ)* .

As we have explained before a C^{1+} hyperbolic diffeomorphism (f, Λ) admits always a marking which is not necessarily unique.

2.9. Leaf n -cylinders and leaf n -gaps. For $\iota = s$ or u , an ι -leaf primary cylinder of a Markov rectangle R is a spanning ι -leaf segment of R . For $n \geq 1$, an ι -leaf n -cylinder of R is an ι -leaf segment I such that

- (i) $f_\iota^n I$ is an ι -leaf primary cylinder of a Markov rectangle M ;
- (ii) $f_\iota^n(\ell^{\iota'}(x, R)) \subset M$ for every $x \in I$.

For $n \geq 2$, an ι -leaf n -gap G of R is an ι -leaf gap $\{x, y\}$ in a Markov rectangle R such that n is the smallest integer such that both leaves $f_\iota^{n-1}\ell^{\iota'}(x, R)$ and $f_\iota^{n-1}\ell^{\iota'}(y, R)$ are contained in ι' -boundaries of Markov rectangles; An ι -leaf primary gap G is the image $f_\iota G'$ by f_ι of an ι -leaf 2-gap G' .

We note that an ι -leaf segment I of a Markov rectangle R can be simultaneously an n_1 -cylinder, $(n_1 + 1)$ -cylinder, $\dots$, n_2 -cylinder of R if $f^{n_1}(I)$, $f^{n_1+1}(I)$, $\dots$, $f^{n_2}(I)$ are all spanning ι -leaf segments. Furthermore, if I is an ι -leaf segment contained in the common boundary of two Markov rectangles R_i and R_j then I can be an n_1 -cylinder of R_i and an n_2 -cylinder of R_j with n_1 distinct of n_2 . If $G = \{x, y\}$ is an ι -gap of R contained in the interior of R then there is a unique n such that G is an n -gap. However, if $G = \{x, y\}$ is contained in the common boundary of two Markov




FIGURE 3. A basic stable holonomy from I to J .

rectangles R_i and R_j then G can be an n_1 -gap of R_i and an n_2 -gap of R_j with n_1 distinct of n_2 . Since the number of Markov rectangles $R_1, \dots, R_k$ is finite, there is $C \geq 1$ such that, in all the above cases for cylinders and gaps we have $|n_2 - n_1| \leq C$.

We say that a leaf segment K is the i -th *mother* of an n -cylinder or an n -gap J of R if $J \subset K$ and K is a leaf $(n - i)$ -cylinder of R . We denote K by $m^i J$.

By the properties of a Markov partition, the smallest full ι -leaf $\hat{K}$ containing a leaf n -cylinder K of a Markov rectangle R is equal to the union of all smallest full ι -leaves containing either a leaf $(n + j)$ -cylinder or a leaf $(n + i)$ -gap of R , with $i \in \{1, \dots, j\}$, contained in K .

2.10. Metric on Λ . We say that a rectangle R is an (n_s, n_u) -rectangle if there is $x \in R$ such that, for $\iota = s$ and u , the spanning leaf segments $\ell^\iota(x, R)$ are either an ι -leaf n_ι -cylinder or the union of two such cylinders with a common endpoint.

The reason for allowing the possibility of the spanning leaf segments being inside two touching cylinders is to allow us to regard geometrically very small rectangles intersecting a common boundary of two Markov rectangles to be small in the sense of having n_s and n_u large.

If $x, y \in \Lambda$ and $x \neq y$ then $d_\Lambda(x, y) = 2^{-n}$ where n is the biggest integer such that both x and y are contained in an (n_s, n_u) -rectangle with $n_s \geq n$ and $n_u \geq n$. Similarly if I and J are ι -leaf segments then $d_\Lambda(I, J) = 2^{-n_\iota}$ where $n_\iota = 1$ and n_ι is the biggest integer such that both I and J are contained in an (n_s, n_u) -rectangle.

2.11. Basic holonomies. Suppose that x and y are two points inside any rectangle R of Λ . Let $\ell(x, R)$ and $\ell(y, R)$ be two stable leaf segments respectively containing x and y and inside R . Then we define $\theta : \ell(x, R) \rightarrow \ell(y, R)$ by $\theta(w) = [w, y]$. Such maps are called the *basic stable holonomies* (see Figure 3). They generate the pseudo-group of all stable holonomies. Similarly we define the basic unstable holonomies.

By Theorem 2.1 in [25], the holonomy $\theta : \ell^\iota(x, R) \rightarrow \ell^\iota(y, R)$ has a $C^{1+\alpha}$ extension to the leaves containing $\ell^\iota(x, R)$ and $\ell^\iota(y, R)$, for some $\alpha > 0$.

2.12. Foliated lamination atlas. In this section when we refer to a C^r object r is allowed to take the values $k + \alpha$ where k is a positive integer and $0 < \alpha \leq 1$. Two ι -leaf charts i and j are C^r compatible if whenever U is an open subset of an ι -leaf segment contained in the domains of i and j then $j \circ i^{-1} : i(U) \rightarrow j(U)$ extends to a C^r diffeomorphism of the real line. Such maps are called *chart overlap maps*. A *bounded C^r ι -lamination atlas* $\mathcal{A}^\iota$ is a set of such charts which (a) cover Λ , (b) are pairwise C^r compatible, and (c) the chart overlap maps are uniformly bounded in the C^r norm.

Let $\mathcal{A}^\iota$ be a bounded $C^{1+\alpha}$ ι -lamination atlas, with $0 < \alpha \leq 1$. If $i : I \rightarrow \mathbb{R}$ is a chart in $\mathcal{A}^\iota$ defined on the leaf segment I and K is a leaf segment in I then we define $|K|_i$ to be the length of the minimal closed interval containing $i(K)$. Since the atlas is bounded, if $j : J \rightarrow \mathbb{R}$ is another chart in $\mathcal{A}^\iota$ defined on the leaf segment J which contains K then the ratio between the lengths $|K|_i$ and $|K|_j$ is universally bounded away from 0 and ∞ . If $K' \subset I \cap J$ is another such segment then

we can define the ratio $r_i(K : K') = |K|_i / |K'|_i$. Although this ratio depends upon i , the ratio is exponentially determined in the sense that if T is the smallest segment containing both K and K' then

$$r_j(K : K') \in (1 \pm \mathcal{O}(|T|_i^\alpha)) r_i(K : K') .$$

This follows from the $C^{1+\alpha}$ smoothness of the overlap maps and Taylor's Theorem.

A C^r lamination atlas $\mathcal{A}^\iota$ has *bounded geometry* (i) if f is a C^r diffeomorphism with C^r norm uniformly bounded in this atlas; (ii) if for all pairs I_1, I_2 of ι -leaf n -cylinders or ι -leaf n -gaps with a common point, we have that $r_i(I_1 : I_2)$ is uniformly bounded away from 0 and ∞ with the bounds being independent of i , I_1 , I_2 and n ; and (iii) for all endpoints x and y of an ι -leaf n -cylinder or ι -leaf n -gap I , we have that $|I|_i \leq \mathcal{O}((d_\Lambda(x, y))^\beta)$ and $d_\Lambda(x, y) \leq \mathcal{O}(|I|_i^\beta)$, for some $0 < \beta < 1$, independent of i , I and n .

A C^r bounded lamination atlas $\mathcal{A}^\iota$ is C^r *foliated* (i) if $\mathcal{A}^\iota$ has bounded geometry; and (ii) if the basic holonomies are C^r and have a C^r norm uniformly bounded in this atlas, except possibly for the dependence upon the rectangles defining the basic holonomy. A bounded lamination atlas $\mathcal{A}^\iota$ is C^{1+} *foliated* if $\mathcal{A}^\iota$ is C^r foliated for some $r > 1$.

2.13. Foliated atlas $\mathcal{A}^\iota(g, \rho)$. Let $g \in \mathcal{T}(f, \Lambda)$ and $\rho = \rho_g$ be a C^{1+} Riemannian metric on the manifold containing Λ . The ι -lamination atlas $\mathcal{A}^\iota(g, \rho)$ determined by ρ is the set of all maps $e : I \rightarrow \mathbb{R}$ where $I = \Lambda \cap \hat{I}$ with $\hat{I}$ a full ι -leaf segment, such that e extends to an isometry between the induced Riemannian metric on $\hat{I}$ and the Euclidean metric on the reals. We call the maps $e \in \mathcal{A}^\iota(\rho)$ the ι -lamination charts. If I is an ι -leaf segment (or a full ι -leaf segment) then by $|I|_\rho$ we mean the length in the Riemannian metric ρ of the minimal full ι -leaf containing I . By Theorem 2.2 in [25], the lamination atlas $\mathcal{A}^\iota(g, \rho)$ is C^{1+} foliated for $\iota = \{s, u\}$.

3. SOLENOID FUNCTIONS

In this section, we construct the stable and unstable solenoid functions, and we prove an equivalence between C^{1+} hyperbolic diffeomorphisms and pairs of stable and unstable solenoid functions.

3.1. HR-Hölder ratios. A *HR-structure* associates an affine structure to each stable and unstable leaf segment in such a way that these vary Hölder continuously with the leaf and are invariant under f .

An affine structure on a stable or unstable leaf is equivalent to a *ratio function* $r(I : J)$ which can be thought of as prescribing the ratio of the size of two leaf segments I and J in the same stable or unstable leaf. A *ratio function* $r(I : J)$ is positive (we recall that each leaf segment has at least two distinct points) and continuous in the endpoints of I and J . Moreover,

$$(3.1) \quad r(I : J) = r(J : I)^{-1} \text{ and } r(I_1 \cup I_2 : K) = r(I_1 : K) + r(I_2 : K)$$

provided I_1 and I_2 intersect at most in one of their endpoints.

We say that r is a ι -ratio function if (i) for all ι -leaf segments K , $r(I : J)$ ($I, J \subset K$) defines a ratio function on K ; (ii) r is invariant under f , i.e. $r(I : J) = r(fI : fJ)$ for all ι -leaf segments; and (iii) for every basic ι -holonomy map $\theta : I \rightarrow J$ between the leaf segment I and the leaf segment J defined with respect to a rectangle R and for every ι -leaf segment $I_0 \subset I$ and every ι -leaf segment or gap $I_1 \subset I$,

$$(3.2) \quad \left| \log \frac{r(\theta I_0 : \theta I_1)}{r(I_0 : I_1)} \right| \leq \mathcal{O}((d_\Lambda(I, J))^\varepsilon)$$

where $\varepsilon \in (0, 1)$ depends upon r and the constant of proportionality also depends upon R , but not on the segments considered.

A *HR-structure* is a pair (r_s, r_u) consisting of a stable and an unstable ratio function.




FIGURE 4. The embedding $e : I \rightarrow \mathbb{R}$.

3.2. Realised ratio functions. Let $(g, \Lambda) \in \mathcal{T}(f, \Lambda)$ and let $\mathcal{A}(g, \rho)$ be an ι -lamination atlas which is C^{1+} foliated. Let $|I| = |I|_\rho$ for every ι -leaf segment I . By hyperbolicity of g in Λ , there are $0 < \nu < 1$ and $C > 0$ such that for all ι -leaf segments I and all $m \geq 0$ we get $|g_{\iota'}^m I| \leq C\nu^m |I|$. Thus, using the mean value theorem and the fact that g_ι is C^r , for all short leaf segments K and all leaf segments I and J contained in it, the ι -realised ratio function $r_{g,\iota}$ given by

$$\begin{aligned} r_{g,\iota}(I : J) &= \lim_{n \rightarrow \infty} \frac{|g_{\iota'}^n I|}{|g_{\iota'}^n J|} \\ &= \frac{|g_{\iota'}^m I|}{|g_{\iota'}^m J|} \prod_{n=m}^{\infty} \left(\frac{|g_{\iota'}^{n+1} I|}{|g_{\iota'}^{n+1} J|} \frac{|g_{\iota'}^n J|}{|g_{\iota'}^n I|} \right) \\ &\in \frac{|g_{\iota'}^m I|}{|g_{\iota'}^m J|} \prod_{n=m}^{\infty} (1 \pm \mathcal{O}(\nu^n |K|^\alpha)) \\ &\subset \frac{|g_{\iota'}^m I|}{|g_{\iota'}^m J|} (1 \pm \mathcal{O}(\nu^n |K|^\alpha)) \end{aligned}$$

is well-defined, where $\alpha = \min\{1, r-1\}$. This construction gives the HR-structure on Λ determined by g . By [24], we get the following equivalence:

Theorem 3.1. *The map $g \rightarrow (r_{g,s}, r_{g,u})$ determines a one-to-one correspondence between C^{1+} conjugacy classes in $\mathcal{T}(f, \Lambda)$ and HR-structures.*

3.3. Lamination atlas. Given an ι -ratio function r , we define the embeddings $e : I \rightarrow \mathbb{R}$ by

$$(3.3) \quad e(x) = r(\ell(\xi, x), \ell(\xi, R))$$

where ξ is an endpoint of the ι -leaf segment I and R is a Markov rectangle containing ξ (see Figure 4). For this definition it is not necessary that R contains I . We denote the set of all these embeddings e by $\mathcal{A}(r)$.

The embeddings e of $\mathcal{A}(r)$ have overlap maps with affine extensions, therefore the atlas $\mathcal{A}(r)$ extends to a $C^{1+\alpha}$ lamination structure $\mathcal{L}(r)$. By Proposition 4.2 in [25], we obtain that $\mathcal{A}(r)$ is a C^{1+} foliated atlas.

Let $g \in \mathcal{T}(f, \Lambda)$ and $\mathcal{A}(g, \rho)$ a C^{1+} foliated ι -lamination atlas determined by a Riemmanian metric ρ . Putting together Proposition 2.5 and Proposition 3.5 of [24], we get that the overlap map $e_1 \circ e_2^{-1}$ between a chart $e_1 \in \mathcal{A}(g, \rho)$ and a chart $e_2 \in \mathcal{A}(r_{g,\iota})$ has a C^{1+} diffeomorphic extension to the reals. Therefore, the atlases $\mathcal{A}(g, \rho)$ and $\mathcal{A}(r_{g,\iota})$ determine the same C^{1+} foliated ι -lamination. In particular, for all short leaf segments K and all leaf segments I and J contained in it, we obtain that

$$r_{g,\iota}(I : J) = \lim_{n \rightarrow \infty} \frac{|g_{\iota'}^n I|_\rho}{|g_{\iota'}^n J|_\rho} = \lim_{n \rightarrow \infty} \frac{|g_{\iota'}^n I|_{i_n}}{|g_{\iota'}^n J|_{i_n}}$$




FIGURE 5. The f -matching condition for ι -leaf segments.

where i_n is any chart in $\mathcal{A}(r_{g,\iota})$ containing the segment $g_{i_n}^n K$ in its domain.

3.4. Realised solenoid functions. For $\iota = s$ and u , let $\mathbf{S}^\iota$ denote the set of all ordered pairs (I, J) of ι -leaf segments with the following properties:

- (i) The intersection of I and J consists of a single endpoint.
- (ii) if $\delta_{f,\iota} = 1$ then I and J are primary ι -leaf cylinders.
- (iii) if $0 < \delta_{f,\iota} < 1$ then $f_\iota I$ is an ι -leaf 2-cylinder of a Markov rectangle R and $f_\iota J$ is an ι -leaf 2-gap also of the same Markov rectangle R .

(See Section 2.6 for the definitions of leaf cylinders and gaps). Pairs (I, J) where both are primary cylinders are called *leaf-leaf pairs*. Pairs (I, J) where J is a gap are called *leaf-gap pairs* and in this case we refer to J as a *primary gap*. The set $\mathbf{S}^\iota$ has a very nice topological structure. If $\delta_{f,\iota} = 1$ then the set $\mathbf{S}^\iota$ is isomorphic to a finite union of intervals, and if $\delta_{f,\iota} < 1$ then the set $\mathbf{S}^\iota$ is isomorphic to an embedded Cantor set.

We define a pseudo-metric $d_{\mathbf{S}^\iota} : \mathbf{S}^\iota \times \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ on the set $\mathbf{S}^\iota$ by

$$d_{\mathbf{S}^\iota}((I, J), (I', J')) = \max \{d_\Lambda(I, I'), d_\Lambda(J, J')\}.$$

Let $g \in \mathcal{T}(f, \Lambda)$. For $\iota = s$ and u , we call the restriction of an ι -ratio function $r_{g,\iota}$ to $\mathbf{S}^i$ a *realised solenoid function* $\sigma_{g,\iota}$. By construction, for $\iota = s$ and u , the restriction of an ι -ratio function to $\mathbf{S}^i$ gives an Hölder continuous function satisfying the matching condition, the boundary condition and the cylinder-gap condition as we now proceed to describe.

3.5. Hölder continuity of solenoid functions. This means that for $t = (I, J)$ and $t' = (I', J')$ in $\mathbf{S}^\iota$, $|\sigma_\iota(t) - \sigma_\iota(t')| \leq \mathcal{O}((d_{\mathbf{S}^\iota}(t, t'))^\alpha)$. The Hölder continuity of $\sigma_{g,\iota}$ and the compactness of its domain imply that $\sigma_{g,\iota}$ is bounded away from zero and infinity.

3.6. Matching condition. Let $(I, J) \in \mathbf{S}^\iota$ be a pair of primary cylinders and suppose that we have pairs

$$(I_0, I_1), (I_1, I_2), \dots, (I_{n-2}, I_{n-1}) \in \mathbf{S}^\iota$$

of primary cylinders such that $f_\iota I = \bigcup_{j=0}^{k-1} I_j$ and $f_\iota J = \bigcup_{j=k}^{n-1} I_j$. Then

$$\frac{|f_\iota I|}{|f_\iota J|} = \frac{\sum_{j=0}^{k-1} |I_j|}{\sum_{j=k}^{n-1} |I_j|} = \frac{1 + \sum_{j=1}^{k-1} \prod_{i=1}^j |I_i|/|I_{i-1}|}{\sum_{j=k}^{n-1} \prod_{i=1}^j |I_i|/|I_{i-1}|}.$$

Hence, noting that $g|_\Lambda = f|_\Lambda$, the realised solenoid function $\sigma_{g,\iota}$ must satisfy the *matching condition* (see Figure 5) for all such leaf segments:

$$(3.4) \quad \sigma_{g,\iota}(I : J) = \frac{1 + \sum_{j=1}^{k-1} \prod_{i=1}^j \sigma_{g,\iota}(I_i : I_{i-1})}{\sum_{j=k}^{n-1} \prod_{i=1}^j \sigma_{g,\iota}(I_i : I_{i-1})}.$$




FIGURE 6. The boundary condition for ι -leaf segments.

3.7. Boundary condition. If the stable and unstable leaf segments have Hausdorff dimension equal to 1, then leaf segments I in the boundaries of Markov rectangles can sometimes be written as the union of primary cylinders in more than one way. This gives rise to the existence of a boundary condition that the realised solenoid functions have to satisfy as we pass to explain.

If J is another leaf segment adjacent to the leaf segment I then the value of $|I|/|J|$ must be the same whichever decomposition we use. If we write $J = I_0 = K_0$ and I as $\bigcup_{i=1}^m I_i$ and $\bigcup_{j=1}^n K_j$ where the I_i and K_j are primary cylinders with $I_i \neq K_j$ for all i and j , then the above two ratios are

$$\sum_{i=1}^m \prod_{j=1}^i \frac{|I_j|}{|I_{j-1}|} = \frac{|I|}{|J|} = \sum_{i=1}^n \prod_{j=1}^i \frac{|K_j|}{|K_{j-1}|}.$$

Thus, noting that $g|\Lambda = f|\Lambda$, a realised solenoid function $\sigma_{g,\iota}$ must satisfy the following *boundary condition* (see Figure 6) for all such leaf segments:

$$(3.5) \quad \sum_{i=1}^m \prod_{j=1}^i \sigma_{g,\iota}(I_j : I_{j-1}) = \sum_{i=1}^n \prod_{j=1}^i \sigma_{g,\iota}(K_j : K_{j-1}).$$

3.8. Scaling function. If the ι -leaf segments have Hausdorff dimension less than one and the ι' -leaf segments have Hausdorff dimension equal to 1, then a primary cylinder I in the ι -boundary of a Markov rectangle can also be written as the union of gaps and cylinders of other Markov rectangles. This gives rise to the existence of a cylinder-gap condition that the ι -realised solenoid functions have to satisfy.

Before defining the cylinder-gap condition, we will introduce the scaling function that will be useful to express the cylinder-gap condition, and also, in Definitions 3.2, the bounded equivalence classes of solenoid functions and, in Definition 8.2, the (δ, P) -bounded solenoid equivalence classes of a Gibbs measure.

Let scl^ι be the set of all pairs (K, J) of ι -leaf segments with the following properties:

- (i) K is a leaf n_1 -cylinder or an n_1 -gap segment for some $n_1 > 1$;
- (ii) J is a leaf n_2 -cylinder or an n_2 -gap segment for some $n_2 > 1$;
- (iii) $m^{n_1-1}K$ and $m^{n_2-1}J$ are the same primary cylinder.

Lemma 3.2. *Every fuction $\sigma_\iota : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ has a canonical extension s_ι to scl^ι . Furthermore, if σ_ι is the restriction of a ratio function $r_\iota|_{\mathbf{S}^\iota}$ to $\mathbf{S}^\iota$ then $s_\iota = r_\iota|_{\text{scl}^\iota}$.*

Remark 3.3. *The above map $s_\iota : \text{scl}^\iota \rightarrow \mathbb{R}^+$ is the scaling function determined by the solenoid function $\sigma_\iota : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$.*

Proof of Lemma 3.2. We are going to give an explicit construction of a realised scaling function $s_{g,\iota}$ from a realised solenoid function $\sigma_{g,\iota}$ with the property that $s_{g,\iota} = r_{g,\iota}|\text{scl}^\iota$ where $r_{g,\iota}$ is a ratio function, i.e for every $(K, J) \in \text{scl}^\iota$ we have

$$s_{g,\iota}(K, J) = \lim_{k \rightarrow \infty} \frac{|g_{\iota'}^k K|_\rho}{|g_{\iota'}^k J|_\rho}$$

where $\mathcal{A}(g, \rho)$ is an ι -lamination atlas.

This construction is canonical and applies to every function $\sigma_\iota : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ determining a canonical extension $s_\iota : \text{scl}^\iota \rightarrow \mathbb{R}^+$ of σ_ι .

Let us proceed to construct the ι -scaling function $s : S^\iota \rightarrow \mathbb{R}^+$ from an ι -solenoid function σ . Suppose that J is an ι -leaf n -cylinder or n -gap. Then there are pairs

$$(I_0, I_1), (I_1, I_2), \dots, (I_{l-1}, I_l) \in \mathbf{S}^\iota$$

such that $mJ = \bigcup_{j=0}^l f_{\iota'}^{n-1} I_j$ and $J = f_{\iota'}^{n-1} I_s$ for some $0 \leq s \leq l$. Let us denote $f_{\iota'}^{n-1} I_j$ by I'_j for every $0 \leq s \leq l$. Then

$$\frac{|mJ|}{|J|} = \sum_{j=0}^l \frac{|I'_j|}{|I'_s|} = 1 + \sum_{j=0}^{s-1} \prod_{i=s}^{j+1} \frac{|I'_{i-1}|}{|I'_i|} + \sum_{j=s+1}^l \prod_{i=s}^{j-1} \frac{|I'_{i+1}|}{|I'_i|},$$

where the first sum above is empty if $s = 0$, and the second sum above is empty if $s = l$. Therefore, we define the extension s from σ to the pairs (mJ, J) by

$$s(mJ, J) = 1 + \sum_{j=0}^{s-1} \prod_{i=s}^{j+1} \sigma(I_{i-1}, I_i) + \sum_{j=s+1}^l \prod_{i=s}^{j-1} \sigma(I_{i+1}, I_i),$$

where the first sum above is empty if $s = 0$, and the second sum above is empty if $s = l$. For every $(K, J) \in \text{scl}^\iota$ there is a primary leaf segment I such that $m^{m_1} K = I = m^{m_2} J$ for some $m_1 \geq 1$ and $m_2 \geq 1$. Then,

$$\frac{|K|}{|J|} = \prod_{j=1}^{m_1} \frac{|m^j J|}{|m^{j-1} J|} \prod_{j=1}^{m_2} \frac{|m^{j-1} K|}{|m^j K|}.$$

Therefore, we define the extension s to (K, J) by

$$s(K, J) = \prod_{j=1}^{m_1} s(m^j J, m^{j-1} J) \prod_{j=1}^{m_2} s(m^{j-1} K, m^j K).$$

Hence, we have constructed a scaling function s from σ on the set scl^ι such that if σ is the restriction of a ratio function $r_{g,\iota}|\mathbf{S}^\iota$ to $\mathbf{S}^\iota$ then $s = r_{g,\iota}|\text{scl}^\iota$. $\square$

3.9. Cylinder-gap condition. Let (I, K) be a leaf-gap pair such that the primary cylinder I is the ι -boundary of a Markov rectangle R_1 . Then the primary cylinder I intersects another Markov rectangle R_2 giving rise to the existence of a cylinder-gap condition that the realised solenoid functions have to satisfy as we proceed to explain. Take the smallest $l \geq 0$ such that $f_{\iota'}^l I \cup f_{\iota'}^l K$ is contained in the intersection of the boundaries of two Markov rectangles M_1 and M_2 . Let M_1 be the Markov rectangle with the property that $M_1 \cap f_{\iota'}^l R_1$ is a rectangle with non-empty interior (and so $M_2 \cap f_{\iota'}^l R_2$ also has non-empty interior). Then, for some positive n , there are distinct n -cylinder and gap leaf segments $J_1, \dots, J_m$ contained in a primary cylinder of M_2 such that $f_{\iota'}^l K = J_m$ and the smallest full ι -leaf segment containing $f_{\iota'}^l I$ is equal to the union $\bigcup_{i=1}^{m-1} \hat{J}_i$, where $\hat{J}_i$ is the smallest




FIGURE 7. The cylinder-gap condition for ι -leaf segments.

full ι -leaf segment containing J_i . Hence,

$$\frac{|f_\iota^l I|}{|f_\iota^l K|} = \sum_{i=1}^{m-1} \frac{|J_i|}{|J_m|}.$$

Hence, noting that $g|\Lambda = f|\Lambda$, a realised solenoid function $\sigma_{g,\iota}$ must satisfy the *cylinder-gap condition* (see Figure 7) for all such leaf segments:

$$\sigma_{g,\iota}(I, K) = \sum_{i=1}^{m-1} s_{g,\iota}(J_i, J_m)$$

where $s_{g,\iota}$ is the scaling function determined by the solenoid function $\sigma_{g,\iota}$.

3.10. Solenoid functions. Now, we are ready to present the definition of an ι -solenoid function.

Definition 3.1. *An Hölder continuous function $\sigma_\iota : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ is an ι -solenoid function if σ_ι satisfies the matching condition, the boundary condition and the cylinder-gap condition.*

We denote by $\mathcal{PS}(f)$ the set of pairs (σ_s, σ_u) of stable and unstable solenoid functions.

Remark 3.4. *Let $\sigma_\iota : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ be an ι -solenoid function. The matching, the boundary and the cylinder-gap conditions are trivially satisfied except in the following cases:*

- (i) *The matching condition if $\delta_{f,\iota} = 1$.*
- (ii) *The boundary condition if $\delta_{f,s} = \delta_{f,u} = 1$.*
- (iii) *The cylinder-gap condition if $\delta_{f,\iota} < 1$ and $\delta_{f,\iota'} = 1$.*

Lemma 3.5. *The map $r_\iota \rightarrow r_\iota|_{\mathbf{S}^\iota}$ gives a one-to-one correspondence between ι -ratio functions and ι -solenoid functions.*

Proof. Every ι -ratio function restricted to the set $\mathbf{S}^\iota$ determines an ι -solenoid function $r_\iota|_{\mathbf{S}^\iota}$. Now we prove the converse. Since the solenoid functions are continuous and their domains are compact they are bounded away from 0 and ∞ . By this boundedness and the f -matching condition of the solenoid functions and by iterating the domains $\mathbf{S}^s$ and $\mathbf{S}^u$ of the solenoid functions backward and forward by f , we determine the ratio functions r^s and r^u at very small (and large) scales, such that f leaves the ratios invariant. Then, using the boundedness again, we extend the ratio functions to all pairs of small adjacent leaf segments by continuity. By the boundary condition and the cylinder-gap condition of the solenoid functions, the ratio functions are well determined at the boundaries of the Markov rectangles. Using the Hölder continuity of the solenoid function, we deduce inequality (3.2).

The set $\mathcal{PS}(f)$ of all pairs (σ_s, σ_u) has a natural metric. Combining Theorem 3.1 with Theorem 3.5, we obtain that the set $\mathcal{PS}(f)$ forms a moduli space for the C^{1+} conjugacy classes of C^{1+} hyperbolic diffeomorphisms $g \in \mathcal{T}(f, \Lambda)$:

Corollary 3.6. *The map $g \rightarrow (r_{g,s}|_{\mathbf{S}^s}, r_{g,u}|_{\mathbf{S}^u})$ determines a one-to-one correspondence between C^{1+} conjugacy classes of $g \in \mathcal{T}(f, \Lambda)$ and pairs of solenoid functions in $\mathcal{PS}(f)$.*

Definition 3.2. We say that any two ι -solenoid functions $\sigma_1 : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ and $\sigma_2 : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ are in the same bounded equivalence class if the corresponding scaling functions $s_1 : \text{scl}^\iota \rightarrow \mathbb{R}^+$ and $s_2 : \text{scl}^\iota \rightarrow \mathbb{R}^+$ satisfy the following property: There is $C > 0$ such that for every ι -leaf $(i+1)$ -cylinder or $(i+1)$ -gap J

$$(3.6) \quad |\log s_1(J, m^i J) - \log s_2(J, m^i J)| < C .$$

Later, in Lemma 9.2, we prove that two C^{1+} hyperbolic diffeomorphisms g_1 and g_2 are Lipschitz conjugate if, and only if, the solenoid functions $s_{g_1, \iota}$ and $s_{g_2, \iota}$ are in the same bounded equivalence class for ι equal to s and u .

4. SELF-RENORMALISABLE STRUCTURES

In this section, we construct the stable and unstable self-renormalisable structures living in 1-dimensional spaces, and we prove an equivalence between C^{1+} hyperbolic diffeomorphisms and pairs of stable and unstable self-renormalisable structures.

4.1. Train-tracks. Roughly speaking train-tracks are the optimal leaf-quotient spaces on which the stable and unstable Markov maps induced by the action of f on leaf segments are local homeomorphisms.

For each Markov rectangle R let t'_R be the set of ι' -segments of R . Thus by the local product structure one can identify t'_R with any spanning ι -leaf segment $\ell^\iota(x, R)$ of R .

We form the space $\mathbf{B}^\iota$ by taking the disjoint union $\bigsqcup_{R \in \mathcal{R}} t'_R$ (union over all Markov rectangles R of the Markov partition $\mathcal{R}$) and identifying two points $I \in t'_R$ and $J \in t'_{R'}$ if either (i) the ι' -leaf segments I and J are ι' -boundaries of Markov rectangles and their intersection contains at least a point which is not an endpoint of I or J or (ii) there is a sequence $I = I_1, \dots, I_n = J$ such that all I_i, I_{i+1} are both identified in the sense of (i). This space is called the ι -train-track and is denoted $\mathbf{B}^\iota$.

Let $\pi_{\mathbf{B}^\iota} : \bigsqcup_{R \in \mathcal{R}} R \rightarrow \mathbf{B}^\iota$ be the natural projection sending $x \in R$ to the point in $\mathbf{B}^\iota$ represented by $\ell^\iota(x, R)$. A *topologically regular point* I in $\mathbf{B}^\iota$ is a point with a unique preimage under $\pi_{\mathbf{B}^\iota}$ (i.e. the pre-image of I is not a union of distinct ι' -boundaries of Markov rectangles). If a point has more than one preimage by $\pi_{\mathbf{B}^\iota}$ then we call it a *junction*. Since there are only a finite number of ι' -boundaries of Markov rectangles there are only finitely many junctions (see Figures 1).

Let d_ι be the metric on $\mathbf{B}^\iota$ defined as follows: if $\xi, \eta \in \mathbf{B}^\iota$, $d_{\mathbf{B}^\iota}(\xi, \eta) = d_\Lambda(\xi, \eta)$.

4.2. Train-track segments and charts. We say that I_T is a *train-track segment* if there is an ι -leaf segment I , not intersecting ι -boundaries of Markov rectangles, such that $\pi_\iota|_I$ is an injection and $\pi_\iota(I) = I_T$. Let $\mathcal{A}$ be an ι -lamination atlas (take for instance $\mathcal{A}$ equal to $\mathcal{A}^\iota(f, \rho)$ or to $\mathcal{A}(r_{f, \iota})$). The chart $i : I \rightarrow \mathbb{R}$ in $\mathcal{A}$ determines a *train-track chart* $i_T : I_T \rightarrow \mathbb{R}$ for I_T given by $i_T = i \circ \pi_\iota^{-1}$. We denote by $\mathcal{B}$ the set of all train-track charts for all train-track segments determined by $\mathcal{A}$.

Two train-track charts (i_T, I_T) and (j_T, J_T) on the train-track $\mathbf{B}^\iota$ are C^{1+} -compatible if the *overlap map* $j_T \circ i_T^{-1} : i_T(I_T \cap J_T) \rightarrow j_T(I_T \cap J_T)$ has a C^{1+} extension. A C^{1+} atlas $\mathcal{B}$ is a set of C^{1+} -compatible charts with the following property: For every short train-track segment K_T there is a chart $(i_T, I_T) \in \mathcal{B}$ such that $K_T \subset I_T$. A C^{1+} *structure* $\mathcal{S}$ on $\mathbf{B}^\iota$ is a maximal set of C^{1+} -compatible charts with a given atlas $\mathcal{B}$ on $\mathbf{B}^\iota$. We say that two C^{1+} structures $\mathcal{S}$ and $\mathcal{S}'$ are in the same *Lipschitz equivalence class* if for every chart in $\mathcal{S}$ and every chart in $\mathcal{S}'$ the overlap map $e_1 \circ e_2^{-1}$ has a bi-Lipschitz extension.

Given any train-track charts $i_T : I_T \rightarrow \mathbb{R}$ and $j_T : J_T \rightarrow \mathbb{R}$ in $\mathcal{B}$, the overlap map $j_T \circ i_T^{-1} : i_T(I_T \cap J_T) \rightarrow j_T(I_T \cap J_T)$ is equal to $j_T \circ i_T^{-1} = j \circ h \circ i^{-1}$ where $i = i_T \circ \pi_\iota : I \rightarrow \mathbb{R}$ and $j = j_T \circ \pi_\iota : J \rightarrow \mathbb{R}$ are charts in $\mathcal{A}$, and

$$h : i^{-1}(i_T(I_T \cap J_T)) \rightarrow j^{-1}(j_T(I_T \cap J_T))$$

is a basic ι -holonomy. Let us denote by $\mathcal{B}^\iota(g, \rho)$ and $\mathcal{B}(r_{g,\iota})$ the train-track atlases determined respectively by $\mathcal{A}^\iota(g, \rho)$ and $\mathcal{A}(r_{g,\iota})$ with $g \in \mathcal{T}(f, \Lambda)$. Since $\mathcal{A}^\iota(g, \rho)$ and $\mathcal{A}(r_{g,\iota})$ are C^{1+} foliated atlases, there is $\eta > 0$ such that, for all train-track charts i_T and j_T in $\mathcal{B}^\iota(g, \rho)$ (or in $\mathcal{B}(r_{g,\iota})$), the overlap maps $j_T \circ i_T^{-1} = j \circ h \circ i^{-1}$ have $C^{1+\eta}$ diffeomorphic extensions with a uniform bound for the $C^{1+\eta}$ norm. Hence, $\mathcal{B}^\iota(g, \rho)$ and $\mathcal{B}(r_{g,\iota})$ are $C^{1+\eta}$ atlas.

4.3. Markov maps. The *Markov map* $\tau_\iota : \mathbf{B}^\iota \rightarrow \mathbf{B}^\iota$ is the mapping induced by the action of f on leaf segments i.e. it is defined as follows: if $I \in \mathbf{B}^\iota$, $\tau_\iota I = \pi_{\mathbf{B}^\iota}(f_\iota I)$ is the ι' -leaf segment containing the f_ι -image of the ι' -leaf segment I (for simplicity of notation we use the same symbols for the Markov maps as for the shift maps). This map τ_ι is a local homeomorphism because f_ι sends a short ι -leaf segment homeomorphically onto a short ι -leaf segment. For simplicity of notation, we will denote τ_ι by f_ι through the paper.

Given a topological chart (e, U) on the train-track $\mathbf{B}^\iota$ and a train-track segment $C \subset U$, we denote by $|C|_e$ the length of the smallest interval containing $e(C)$. We say that f_ι has *bounded geometry* in a C^{1+} atlas $\mathcal{B}$ if there is $\kappa_1 > 0$ such that, for every n -cylinder C_1 and n -cylinder or n -gap C_2 with a common endpoint with C_1 , we have $\kappa_1^{-1} < |C_1|_e/|C_2|_e < \kappa_1$, where the lengths are measured in any chart (e, U) of the atlas such that $C_1 \cup C_2 \subset U$. Hence there is $\kappa_2 > 0$ and $0 < \nu < 1$ such that $|C|_e \leq \kappa_2 \nu^n$ for every n -cylinder or n -gap C . This property is equivalent to the Markov map f_ι being uniformly expanding in $\mathbf{B}^\iota$.

Since f on Λ along leaves has affine extensions with respect to the charts in $\mathcal{A}(r^\iota)$ and the basic ι -holonomies have $C^{1+\eta}$ extensions we get that the Markov maps τ_ι also have $C^{1+\eta}$ extensions with respect to the charts in $\mathcal{B}(r^\iota)$ for some $\eta > 0$. Since $\mathcal{A}(r^\iota)$ has bounded geometry, we obtain that f_ι also has *bounded geometry* in $\mathcal{B}(r^\iota)$. Since, for every $g \in \mathcal{T}(f, \Lambda)$, the C^{1+} lamination atlas $\mathcal{A}(g, \rho_g)$ has bounded geometry we obtain that the Markov map f_ι has $C^{1+\eta}$ extensions with respect to the charts in $\mathcal{B}(g, \rho_g)$, for some $\eta > 0$, and has bounded geometry.

4.4. Holonomy pseudo-groups on $\mathbf{B}^\iota$. The elements $\theta_\iota = \theta_{f_\iota}$ of the *holonomy pseudo-group* on $\mathbf{B}^\iota$ are the mappings defined as follows. Suppose that I and J are ι -leaf segments and $h : I \rightarrow J$ a holonomy. Then it follows from the definition of the train-track $\mathbf{B}^\iota$ that the map $\theta : \pi_{\mathbf{B}^\iota}(I) \rightarrow \pi_{\mathbf{B}^\iota}(J)$ given by $\theta(\pi_{\mathbf{B}^\iota}(x)) = \pi_{\mathbf{B}^\iota}(h(x))$ is well-defined. The collection of all such local mappings forms the *basic holonomy pseudo-group* of $\mathbf{B}^\iota$. Note that if x is a junction of $\mathbf{B}^\iota$ then there may be segments I and J containing x such that $I \cap J = \{x\}$. In this case the image of I and J under the holonomies will not agree in that they will map x differently.

4.5. Markings on train-tracks. For $\iota = s$ and u , the Markov partition $\mathcal{R} = \{R_1, \dots, R_m\}$ for (f, Λ) induces a Markov partition $\mathcal{R}^\iota = \{R_1^\iota, \dots, R_m^\iota\}$ for the Markov map $\tau = \tau_\iota$ on the train-track $\mathbf{B}^\iota$. The marking $i : \Theta \rightarrow \Lambda$ determines unique *markings* $i_u : \Theta^u \rightarrow \mathbf{B}^u$ and $i_s : \Theta^s \rightarrow \mathbf{B}^s$ such that $i_u(w_0 w_1 \dots) = \cap_{i \geq 0} R_{w_i}^u$ and $i_s(\dots w_{-1} w_0) = \cap_{i \geq 0} R_{w_i}^u$. We note that $\pi_{\mathbf{B}^\iota} \circ i = i_\iota \circ \pi_\iota$. The map i_ι is continuous, onto $\mathbf{B}^\iota$ and semiconjugates the shift map on Θ^ι to the Markov map on $\mathbf{B}^\iota$. Defining $\varepsilon, \varepsilon' \in \Theta^\iota$ to be equivalent ($\varepsilon \sim \varepsilon'$) if the point $i^\iota(\varepsilon) = i^\iota(\varepsilon')$, we get that the space $\Theta^\iota / \sim$ is homeomorphic to the train-track $\mathbf{B}^\iota$.

Consider the Markov map f_ι on $\mathbf{B}^\iota$ induced by the action of f on ι' -leaves and described above. For $n \geq 1$, an n -cylinder is the projection into $\mathbf{B}^\iota$ of an ι -leaf n -cylinder segment in Λ . Thus, each Markov rectangle in Λ projects in an unique primary ι -leaf segment in $\mathbf{B}^\iota$. For $n \geq 1$, an n -gap of f_ι is the projection into $\mathbf{B}^\iota$ of a ι -leaf n -gap in Λ .

We say that $\mathbf{B}^\iota$ is a *no-gap train-track* if $\mathbf{B}^\iota$ does not have gaps. Otherwise, we call $\mathbf{B}^\iota$ a *gap train-track*.

4.6. Self-renormalisable structures. The C^{1+} structure $\mathcal{S}_\iota$ on $\mathbf{B}^\iota$ is an ι *self-renormalisable* if it has the following properties:




FIGURE 8. A Markov partition for the Smale-shoe f into two rectangles A and B . A representation of the Markov maps $m_s : \Theta^s \rightarrow \Theta^s$ and $m_u : \Theta^u \rightarrow \Theta^u$ for Smale horseshoes.

- (i) in this structure the Markov mapping m_i is a local diffeomorphism and has bounded geometry in some C^{1+} atlas of this structure; and
- (ii) the elements of the basic holonomy pseudo-group are local diffeomorphisms in $\mathcal{S}_i$.

We say that $\mathcal{B}$ is a C^{1+} *self-renormalisable atlas* if $\mathcal{B}$ has bounded geometry and extends to a C^{1+} self-renormalisable structure. By definition, a C^{1+} self-renormalisable structure contains a C^{1+} self-renormalisable atlas.

A C^{1+} foliated ι -lamination atlas $\mathcal{A}$ induces a C^{1+} ι self-renormalisable atlas $\mathcal{B}$ on $\mathbf{B}^\iota$ (and vice-versa) as follows: The holonomies are C^{1+} with respect to the atlas $\mathcal{A}$ and so the charts in $\mathcal{B}$ are C^{1+} compatible, and the basic holonomy pseudo-group of $\mathbf{B}^\iota$ are local diffeomorphisms. Since $\mathcal{A}$ has bounded geometry the Markov mapping τ_i is a local diffeomorphism and also has bounded geometry in $\mathcal{B}$. Therefore, $\mathcal{B}$ is a C^{1+} self-renormalisable atlas and extends to a C^{1+} self-renormalisable structure $\mathcal{S}(\mathcal{B})$ on $\mathbf{B}^\iota$. Since $\mathcal{A}(r_\iota)$ and $\mathcal{A}^\iota(g, \rho_g)$ are C^{1+} foliated ι -lamination atlas we obtain that the atlases $\mathcal{B}(r_\iota)$ and $\mathcal{B}^\iota(g, \rho_g)$ determine respectively C^{1+} self-renormalisable structures $\mathcal{S}(r_\iota)$ and $\mathcal{S}(g, \iota)$.




FIGURE 9. A representation of the Markov maps $m_s : \Theta^s \rightarrow \Theta^s$ and $m_u : \Theta^u \rightarrow \Theta^u$ as maps of the interval for Anosov diffeomorphisms.

Lemma 4.1. *The map $r_\iota \rightarrow \mathcal{S}(r_\iota)$ determines a one-to-one correspondence between ι -ratio functions (or equivalently, ι -solenoid functions $r_\iota|_{\mathbf{B}^\iota}$) and C^{1+} self-renormalisable structures on $\mathbf{B}^\iota$.*

Proof. Every ratio function r_ι determines an unique C^{1+} self-renormalisable $\mathcal{S}$. Conversely, let us prove that a given C^{1+} self-renormalisable structure $\mathcal{S}$ on $\mathbf{B}^\iota$ also determines an unique ratio function $r_{\mathcal{S},\iota}$. Let $\mathcal{B}$ be a bounded atlas for $\mathcal{S}$. Consider a small leaf segment K and two leaf segments I and J contained in K . Since the elements of the basic holonomy pseudo-group on $\mathbf{B}^\iota$ are C^{1+} and the Markov map is also C^{1+} and has bounded geometry we obtain by Taylor's Theorem that the following limit exists

$$(4.1) \quad \begin{aligned} r_{\mathcal{S},\iota}(I : J) &= \lim_{n \rightarrow \infty} \frac{|\pi_\iota f_{\iota'}^n I|_{i_n}}{|\pi_\iota f_{\iota'}^n J|_{i_n}} \\ &\in \frac{|\pi_\iota I|_{i_0}}{|\pi_\iota J|_{i_0}} (1 \pm \mathcal{O}(|\pi_\iota K|_{i_0}^\gamma)) , \end{aligned}$$

where the size of the leaf segments are measured in charts of the bounded atlas $\mathcal{B}$. Furthermore, by [24] and (4.1), the charts in $\mathcal{B}(r_\iota)$ and the charts in $\mathcal{B}$ are C^{1+} equivalent and so determine the same C^{1+} self-renormalisable structure.

4.7. Hyperbolic diffeomorphisms. Let $g \in \mathcal{T}(f, \Lambda)$ and $\mathcal{A}(g, \rho_g)$ be the C^{1+} foliated ι -lamination atlas determined by the Riemannian metric ρ_g . As shown in Section 4.6, the atlas $\mathcal{A}(g, \rho_g)$ induces a C^{1+} self-renormalisable atlas $\mathcal{B}(g, \rho_g)$ on $\mathbf{B}^\iota$ which generates a C^{1+} self-renormalisable structure $\mathcal{S}(g, \iota)$.

Lemma 4.2. *The mapping $g \rightarrow (\mathcal{S}(g, s), \mathcal{S}(g, u))$ gives a 1-1 correspondence between C^{1+} conjugacy classes in $\mathcal{T}(f, \Lambda)$ and pairs $(\mathcal{S}(g, s), \mathcal{S}(g, u))$ of C^{1+} self-renormalisable structures. Furthermore, $r_{g,s} = r_{\mathcal{S}(g,s),s}$ and $r_{g,u} = r_{\mathcal{S}(g,u),u}$.*

Proof. By Lemma 4.1, the pair $(\mathcal{S}_s, \mathcal{S}_u)$ determines a pair $(r_{s,\mathcal{S}}|_{\mathbf{S}^s}, r_{u,\mathcal{S}}|_{\mathbf{S}^u})$ of solenoid functions and vice-versa. By Corollary 3.6, the pair $(r_{s,\mathcal{S}}|_{\mathbf{S}^s}, r_{u,\mathcal{S}}|_{\mathbf{S}^u})$ determines an unique C^{1+} conjugacy class of diffeomorphisms $g \in \mathcal{T}(f, \Lambda)$ which realise the pair $(r_{s,\mathcal{S}}|_{\mathbf{S}^s}, r_{u,\mathcal{S}}|_{\mathbf{S}^u})$ and vice-versa (and so $(\mathcal{S}(g, s), \mathcal{S}(g, u)) = (\mathcal{S}_s, \mathcal{S}_u)$). Furthermore, by Lemma 3.5, we get $r_{g,s} = r_{\mathcal{S}(g,s),s}$ and $r_{g,u} = r_{\mathcal{S}(g,u),u}$.

5. MEASURE SOLENOID FUNCTIONS

In this section, we introduce the following new concepts: stable and unstable measure solenoid functions and stable and unstable measure ratio functions. Later, we will use the measure solenoid functions and the measure ratio functions to determine which Gibbs measures are C^{1+} -realisable by C^{1+} hyperbolic diffeomorphisms and by C^{1+} self-renormalisable structures.

5.1. Gibbs measures. Let us give the definition of an infinite two-sided subshift of finite type $\Theta = \Theta(A)$. The elements of Θ are all infinite two-sided words $w = \dots w_{-1}w_0w_1\dots$ in the symbols $1, \dots, k$ such that $A_{w_iw_{i+1}} = 1$, for all $i \in \mathbb{Z}$. Here $A = (A_{ij})$ is any matrix with entries 0 and 1 such that A^n has all entries positive for some $n \geq 1$. We write $w \stackrel{n_1, n_2}{\sim} w'$ if $w_j = w'_j$ for every $j = -n_1, \dots, n_2$. The metric d on Θ is given by $d(w, w') = 2^{-n}$ if $n \geq 0$ is the largest such that $w \stackrel{n, n}{\sim} w'$. Together with this metric Θ is a compact metric space. The two-sided shift map $\tau : \Theta \rightarrow \Theta$ is the mapping which sends $w = \dots w_{-1}w_0w_1\dots$ to $v = \dots v_{-1}v_0v_1\dots$ where $v_j = w_{j+1}$ for every $j \in \mathbb{Z}$. An (n_1, n_2) -cylinder $\Theta_{w_{-n_1} \dots w_{n_2}}$, where $w \in \Theta$, consists of all those words w' in Θ such that $w \stackrel{n_1, n_2}{\sim} w'$. Let Θ^u be the set of all right-handed words $w_0w_1\dots$ which extend to words $\dots w_0w_1\dots$ in Θ , and, similarly, let Θ^s be the set of all left-handed words $\dots w_{-1}w_0$ which extend

to words $\dots w_{-1}w_0\dots$ in Θ . Then $\pi_u : \Theta \rightarrow \Theta^u$ and $\pi_s : \Theta \rightarrow \Theta^s$ are the natural projection given, respectively, by

$$\pi_u(\dots w_{-1}w_0w_1\dots) = w_0w_1\dots \text{ and } \pi_s(\dots w_{-1}w_0w_1\dots) = \dots w_{-1}w_0.$$

An n -cylinder $\Theta_{w_0\dots w_{n-1}}^u$ is equal to $\pi_u(\Theta_{w_0\dots w_{n-1}})$ and an n -cylinder $\Theta_{w_{-(n-1)}\dots w_0}^s$ is equal to $\pi_s(\Theta_{w_{-(n-1)}\dots w_0})$. Let $\tau_u : \Theta^u \rightarrow \Theta^u$ and $\tau_s : \Theta^s \rightarrow \Theta^s$ be the corresponding one-sided shifts.

Definition 5.1. For $\iota = s$ and u , we say that $s_\iota : \Theta^\iota \rightarrow \mathbb{R}^+$ is an ι -measure scaling function if s_ι is a Hölder continuous function, and for every $\xi \in \Theta^\iota$

$$\sum_{\tau_\iota \eta = \xi} s_\iota(\eta) = 1,$$

where the sum is upon all $\xi \in \Theta^\iota$ such that $\tau_\iota \eta = \xi$.

For $\iota \in \{s, u\}$, a τ -invariant measure ν on Θ determines a unique τ_ι -invariant measure $\nu_\iota = (\pi_\iota)_* \nu$ on Θ^ι . We note that a τ_ι -invariant measure ν_ι on Θ^ι has an unique τ -invariant natural extension to an invariant measure ν on Θ such that $\nu(\Theta_{w_0\dots w_{n_2}}) = \nu_\iota(\Theta_{w_0\dots w_{n_2}}^\iota)$.

Definition 5.2. A τ -invariant measure ν on Θ is a Gibbs measure:

(i) if the function $s_{\nu,u} : \Theta^u \rightarrow \mathbb{R}^+$ given by

$$s_{\nu,u}(w_0w_1\dots) = \lim_{n \rightarrow \infty} \frac{\nu(\Theta_{w_0\dots w_n})}{\nu(\Theta_{w_1\dots w_n})},$$

is well-defined and it is an u -measure scaling function; and

(ii) if the function $s_{\nu,s} : \Theta^s \rightarrow \mathbb{R}^+$ given by

$$s_{\nu,s}(\dots w_1w_0) = \lim_{n \rightarrow \infty} \frac{\nu(\Theta_{w_n\dots w_0})}{\nu(\Theta_{w_n\dots w_1})},$$

is well-defined and it is a s -measure scaling function.

The following theorem follows from the results proved in [23]. It can also be deduced from standard results about Gibbs states such as those in [2].

Theorem 5.1. (Moduli space for Gibbs measures) Let $s_\iota : \Theta^\iota \rightarrow \mathbb{R}^+$ be an ι -measure scaling function for $\iota = s$ or u . Then there is an unique τ -invariant Gibbs measure ν such that $s_{\nu,\iota} = s_\iota$.

5.2. Extended measure scaling function. To present a classification of Gibbs measures C^{1+} -Hausdorff which are realisable by codimension one attractors, we have to define the cylinder-cylinder condition. We will express the cylinder-cylinder condition, in Section 5.3, using the extended measure scaling functions. These extended measure scaling functions are also used to present, in Section 5.2, the δ_ι -bounded solenoid equivalence class of a Gibbs measure.

Throughout the paper, if $\xi \in \Theta^{\iota'}$, we denote by ξ_Λ the leaf primary cylinder segment $i(\pi_{\iota'}^{-1}\xi) \subset \Lambda$. Similarly, if C is an n_ι -cylinder of Θ^ι then we denote by C_Λ the $(1, n_\iota)$ -rectangle $i(\pi_\iota^{-1}C) \subset \Lambda$.

We say that a (n_1, n_2) -cylinder $\theta_{w_{-n_1}\dots w_{n_2}}$ of Θ is an u -symbolic leaf n_2 -cylinder if $n_1 = -\infty$. Similarly, we say that a (n_1, n_2) -cylinder $\theta_{w_{-n_1}\dots w_{n_2}}$ is a s -symbolic leaf n_1 -cylinder if $n_2 = +\infty$. Let $\xi \in \Theta^{\iota'}$ and C be a n -cylinder of Θ^ι . We say that the pair (ξ, C) is ι -admissible if the set

$$\xi.C = \pi_{\iota'}^{-1}C \cap \pi_{\iota'}^{-1}\xi$$

is non-empty (see Figure 10). We note that if the pair (ξ, C) is ι -admissible then $\xi.C$ is an ι -symbolic leaf n -cylinder, and, conversely, any ι -symbolic leaf n -cylinder can be expressed as $\xi.C$ where the pair (ξ, C) is ι -admissible. The set of all ι -admissible pairs (ξ, C) is the ι -measure scaling set msc^ι .




FIGURE 10. An ι -admissible pair (ξ, C) where $\xi_\Lambda = i(\pi_{\iota'}^{-1}\xi)$ and $C_\Lambda = i(\pi_\iota^{-1}C)$.




FIGURE 11. The $(n-j+1)$ -cylinder leaf segment $I = \xi_\Lambda \cap D_\Lambda$ and the primary leaf segment $f^{n-j}(I) = i(\pi_{\iota'}\tau_\iota^{n-j}(\xi.D))$, where $D = m_\iota^{j-1}C$.

Let $C^u = \Theta_{w_0 \dots w_{n-1}}^u$ and $C^s = \Theta_{w_{-(n-1)} \dots w_0}^s$ be the n -cylinders of Θ^u and of Θ^s , respectively. For $i < n$, we denote by $m^i C^u$ the i -th mother $\Theta_{w_0 \dots w_{(n-i-1)}}^u$ of C^u . Similarly, we denote by $m^i C^s$ the i -th mother $\Theta_{w_{-(n-i-1)} \dots w_0}^s$ of C^s .

Given an ι -measure scaling function s , we construct the *extended ι -measure scaling function* $\rho : \text{msc}_\iota \rightarrow \mathbb{R}^+$ of s as follows: If C is a 1-cylinder then we define $\rho_\xi(C) = 1$. If C is a n -cylinder with $n \geq 2$, then we define

$$\rho_\xi(C) = \prod_{j=1}^{n-1} s(\pi_{\iota'}\tau_\iota^{n-j}(\xi.m_\iota^{j-1}C))$$

(see Figure 11). We note that, if $s_{\nu,s}$ is the stable measure scaling function of a Gibbs measure ν , then

$$\rho_\xi(C) = \lim_{m \rightarrow \infty} \frac{\nu(\pi_s \circ \tau^m(\xi.C))}{\nu(\pi_s \circ \tau^m(\xi))}.$$

The unstable case is similar to the one above by taking $-m$ instead of m . Hence, $\rho_\xi(C)$ is the ratio between the measure of $\xi.C$ and the measure of ξ with respect to the conditional measure determined by the Gibbs measure ν in ξ .

Recall that a τ -invariant measure ν on Θ determines a unique τ_u -invariant measure $\nu_u = (\pi_u)_*\nu$ on Θ^u and a unique τ_s -invariant measure $\nu_s = (\pi_s)_*\nu$ on Θ^s . The following theorem follows from [23].

Theorem 5.2. (*Ratio decomposition of a Gibbs measure*) Let $\rho : \text{msc}_\iota \rightarrow \mathbb{R}^+$ be an extended ι -measure scaling function and ν the corresponding τ -invariant Gibbs measure. If C is an n -cylinder




FIGURE 12. The cylinder-cylinder condition for ι -leaf segments.

of Θ^ι then

$$\nu_\iota(C) = \int_{\xi \in M} \rho_\xi(C) \nu_{\iota'}(d\xi).$$

where $M = \pi_{\iota'} \circ \pi_\iota^{-1} C$ a 1-cylinder of $\Theta^{\iota'}$. Furthermore, the ratios $\nu_\iota(C)/\rho_\xi(C)$ are uniformly bounded away from 0 and ∞ .

5.3. Cylinder-cylinder condition. We introduce the cylinder-cylinder condition that we will use to classify all Gibbs measures that are C^{1+} -Hausdorff realizable by codimension one attractors.

Similarly to the cylinder-gap condition given in Section 3.9 for a given solenoid function, we are going to construct the cylinder-cylinder condition for a given measure solenoid function $\sigma_{\nu,\iota}$. Let $\delta_\iota < 1$ and $\delta_{\iota'} = 1$. Let $(I, J) \in \text{Msol}^\iota$ be such that the ι -leaf segment $f_{\iota'} I \cup f_{\iota'} J$ is contained in an ι -boundary K of a Markov rectangle R_1 . Then $f_{\iota'} I \cup f_{\iota'} J$ intersects another Markov rectangle R_2 . Take the smallest $k \geq 0$ such that $f_{\iota'}^k I \cup f_{\iota'}^k J$ is contained in the intersection of the boundaries of two Markov rectangles M_1 and M_2 . Let M_1 be the Markov rectangle with the property that $M_1 \cap f_{\iota'}^k R_1$ is a rectangle with non empty interior, and so $M_2 \cap f_{\iota'}^k R_2$ has also non-empty interior. Then, for some positive n , there are distinct ι -leaf n -cylinders $J_1, \dots, J_m$ contained in a primary cylinder L of M_2 such that $f_{\iota'}^k I = \bigcup_{i=1}^{p-1} J_i$ and $f_{\iota'}^k J = \bigcup_{i=p}^m J_i$. Let $\eta \in \Theta^{\iota'}$ be such that $\eta_\Lambda = L$ and, for every $i = 1, \dots, m$, let D_i be a cylinder of Θ^ι such that $i(\eta.D_i) = J_i$. Let $\xi \in \Theta^{\iota'}$ be such that $\xi_\Lambda = K$ and C_1 and C_2 cylinders of Θ^ι such that $i(\xi.C_1) = f_{\iota'} I$ and $i(\xi.C_2) = f_{\iota'} J$. We say that an ι -extended scaling function ρ satisfies the *cylinder-cylinder condition* (see Figure 12) if for all such leaf segments:

$$\frac{\rho_\xi(C_2)}{\rho_\xi(C_1)} = \frac{\sum_{i=p}^m \rho_\eta(D_i)}{\sum_{i=1}^{p-1} \rho_\eta(D_i)}.$$

5.4. Measure solenoid functions. Let Msol^ι be the set of all pairs (I, J) with the following properties: (a) If $\delta_\iota = 1$ then $\text{Msol}^\iota = \mathbf{S}^\iota$. (b) If $\delta_\iota < 1$ then $f_{\iota'} I$ and $f_{\iota'} J$ are ι -leaf 2-cylinders of a Markov rectangle R such that $f_{\iota'} I \cup f_{\iota'} J$ is an ι -leaf segment, i.e. there is a unique ι -leaf 2-gap between them. Let msol^ι be the set of all pairs $(I, J) \in \text{Msol}^\iota$ such that the leaf segments I and J are not contained in an ι -global leaf containing an ι -boundary of a Markov rectangle. By construction, the set msol^ι is dense in Msol^ι , and for every pair $(C, D) \in \text{msol}^\iota$ there is a unique $\psi \in \Theta^{\iota'}$ and a unique $\xi \in \Theta^{\iota'}$ such that $i(\pi_{\iota'}^{-1}(\psi)) = C$ and $i(\pi_{\iota'}^{-1}(\xi)) = D$. We will denote, in what follows, $i(\pi_{\iota'}^{-1}(\psi))$ by ψ_Λ and $i(\pi_{\iota'}^{-1}(\xi))$ by ξ_Λ .

Lemma 5.3. *Let ν be a Gibbs measure on Θ . The s -measure solenoid function $\sigma_{\nu,s} : \text{msol}^s \rightarrow \mathbb{R}^+$ of ν and the u -measure solenoid function $\sigma_{\nu,u} : \text{msol}^u \rightarrow \mathbb{R}^+$ of ν given by*

$$\sigma_{\nu,s}(\psi_\Lambda, \xi_\Lambda) = \lim_{n \rightarrow \infty} \frac{\nu(\Theta_{\psi_0 \dots \psi_n})}{\nu(\Theta_{\xi_0 \dots \xi_n})}$$

and

$$\sigma_{\nu,u}(\psi_\Lambda, \xi_\Lambda) = \lim_{n \rightarrow \infty} \frac{\nu(\Theta_{\psi_n \dots \psi_0})}{\nu(\Theta_{\xi_n \dots \xi_0})}$$

are both well-defined.

Proof. Let $(I, J) \in \text{msol}^\iota$. By Property (iii) of msol^ι , there is $k = k(I, J)$ such that $f_\iota^k I$ and $f_\iota^k J$ are cylinders with the same mother $m f_\iota^k I = m f_\iota^k J$. Let $(\xi : C)$ and $(\xi : D)$ be the admissible pairs in msc_ι such that $i(\xi.C) = f_\iota^k I$ and $i(\xi.D) = f_\iota^k J$. Since the measure ν is τ -invariant, we obtain that

$$\sigma_{\nu,\iota}(I, J) = \rho_\xi(C) \rho_\xi(D)^{-1},$$

where ρ is the extended scaling function determined by the Gibbs measure ν . Therefore, the ι -measure solenoid function $\sigma_{\nu,\iota}$ is well-defined for $\iota \in \{s, u\}$.

Lemma 5.4. *Suppose $\delta_{f,\iota} = 1$. If an ι -measure solenoid function $\sigma_{\nu,\iota} : \text{msol}^\iota \rightarrow \mathbb{R}^+$ has a continuous extension to $\mathbf{S}^\iota$ then its extension satisfies the matching condition.*

Proof. Let $(J_0, J_1) \in \mathbf{S}^\iota$ be a pair of primary cylinders and suppose that we have pairs

$$(I_0, I_1), (I_1, I_2), \dots, (I_{n-2}, I_{n-1}) \in \mathbf{S}^\iota$$

of primary cylinders such that $f_\iota J_0 = \bigcup_{j=0}^{k-1} I_j$ and $f_\iota J_1 = \bigcup_{j=k}^{n-1} I_j$. Since the set msol^ι is dense in $\mathbf{S}^\iota$ there are pairs $(J_0^l, J_1^l) \in \text{msol}^\iota$ and pairs (I_j^l, I_{j+1}^l) with the following properties:

- (i) $f_\iota J_0^l = \bigcup_{j=0}^{k-1} I_j^l$ and $f_\iota J_1^l = \bigcup_{j=k}^{n-1} I_j^l$.
- (ii) The pair (J_0^l, J_1^l) converges to (J_0, J_1) when l tends to infinity.

Therefore, for every $j = 0, \dots, n-2$ the pair (I_j^l, I_{j+1}^l) converges to (I_j, I_{j+1}) when l tends to infinity. Since ν is a τ -invariant measure, we get that the matching condition

$$\sigma_{\nu,\iota}(J_0^l : J_1^l) = \frac{1 + \sum_{j=1}^{k-1} \prod_{i=1}^j \sigma_{\nu,\iota}(I_j^l : I_{i-1}^l)}{\sum_{j=k}^{n-1} \prod_{i=1}^j \sigma_{\nu,\iota}(I_j^l : I_{i-1}^l)}$$

is satisfied for every $l \geq 1$. Since the extension of $\sigma_{\nu,\iota} : \text{msol}^\iota \rightarrow \mathbb{R}^+$ to the set $\mathbf{S}^\iota$ is continuous, we get that the matching condition also holds for the pairs (J_0, J_1) and $(I_0, I_1), \dots, (I_{n-2}, I_{n-1})$.

Remark 5.5. *We say that an ι -measure solenoid function $\sigma_{\nu,\iota}$ of a Gibbs measure ν satisfies the cylinder-cylinder condition if the extended scaling function of the Gibbs measure ν satisfies the cylinder-cylinder condition.*

5.5. Measure ratio functions. We say that ρ is a ι -measure ratio function if

- (i) $\rho(I : J)$ is well-defined for every pair of ι -leaf segments I and J such that (a) there is an ι -leaf segment K such that $I, J \subset K$, and (b) I or J has non-empty interior;
- (ii) if I is an ι -leaf gap then $\rho(I : J) = 0$ (and $\rho(J : I) = +\infty$);
- (iii) if I and J have non-empty interiors then $\rho(I : J)$ is strictly positive;
- (iv) $\rho(I : J) = \rho(J : I)^{-1}$;
- (v) if I_1 and I_2 intersect at most in one of their endpoints then $\rho(I_1 \cup I_2 : K) = \rho(I_1 : K) + \rho(I_2 : K)$;
- (vi) ρ is invariant under f , i.e. $\rho(I : J) = \rho(fI : fJ)$ for all ι -leaf segments;
- (vii) for every basic ι -holonomy map $\theta : I \rightarrow J$ between the leaf segment I and the leaf segment J defined with respect to a rectangle R and for every ι -leaf segment $I_0 \subset I$ and every ι -leaf segment or gap $I_1 \subset I$,

$$(5.1) \quad \left| \log \frac{\rho(\theta I_0 : \theta I_1)}{\rho(I_0 : I_1)} \right| \leq \mathcal{O}((d_\Lambda(I, J))^\varepsilon)$$

where $\varepsilon \in (0, 1)$ depends upon ρ and the constant of proportionality also depends upon R , but not on the segments considered.

We note that if $\mathbf{B}^\iota$ is a no-gap train-track then an ι -measure ratio function is an ι -ratio function.

Remark 5.6. *A function $\sigma : \text{msol}^\iota \rightarrow \mathbb{R}^+$ that has an Hölder continuous extension to Msol^ι determines a unique extended scaling function ρ , and so we say that σ satisfies the cylinder-cylinder condition if the extended scaling function ρ satisfies the cylinder-cylinder condition.*

Let SOL^ι be the space of all Hölder continuous functions $\sigma_\iota : \text{Msol}^\iota \rightarrow \mathbb{R}^+$ with the following properties:

- (i) If $\mathbf{B}^\iota$ is a no-gap train-track then σ_ι is an ι -solenoid function.
- (ii) If $\mathbf{B}^\iota$ is a gap train-track and $\mathbf{B}^{\iota'}$ is a no-gap train-track then σ_ι satisfies the cylinder-cylinder condition.
- (iii) If $\mathbf{B}^\iota$ and $\mathbf{B}^{\iota'}$ are no-gap train-tracks then σ_ι does not have to satisfy any extra property.

Lemma 5.7. *The map $\rho \rightarrow \rho|_{\text{Msol}^\iota}$ determines an one-to-one correspondence between ι -measure ratio functions and functions contained in SOL^ι .*

Proof. The proof follows similarly to the proof of Lemma 3.5.

Remark 5.8. (i) *By Lemma 5.7, a Gibbs measure ν with an ι -measure solenoid function with an extension $\hat{\sigma}$ to Msol^ι such that $\hat{\sigma} \in SOL^\iota$ determines a unique ι -measure ratio function ρ_ν .*
(ii) *A measure ratio function ρ determines naturally a measure scaling function, and so, by Lemma 5.1, a Gibbs measure ν_ρ .*
(iii) *By Lemma 5.7, a function $\sigma : \text{msol}^\iota \rightarrow \mathbb{R}^+$ with an extension $\hat{\sigma}$ to Msol^ι such that $\hat{\sigma} \in SOL^\iota$ determines an ι -measure ratio function, and, by (ii), a unique Gibbs measure ν such that $\sigma = \sigma_\nu$.*

6. NATURAL GEOMETRIC MEASURES

In this section, we define the natural geometric measures $\mu_{\mathcal{S},\delta}$ associated with a self-renormalisable structure $\mathcal{S}$ and $\delta > 0$. The natural geometric measures are measures determined by the length scaling structure of the cylinders. We will prove that every natural geometric measure is a pushforward of a Gibbs measure with the property that the measure solenoid function determines a measure ratio function. In Section 8, we will show that a Gibbs measure with the property that its measure solenoid function determines a measure ratio function is C^{1+} -realisable by a self-renormalisable structure.

Definition 6.1. *Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on $\mathbf{B}^\iota$. If $\mathbf{B}^\iota$ is a gap train-track let $0 < \delta < 1$, and if $\mathbf{B}^\iota$ is a no-gap train-track let $\delta = 1$.*

- (i) *We say that $\mathcal{S}$ has a natural geometric measure $\mu_\iota = \mu_{\mathcal{S},\delta}$ with pressure $P = P(\mathcal{S},\delta)$ if (a) μ_ι is a f_ι -invariant measure; (b), there exists $\kappa > 1$ such that for all $n \geq 1$ and all n -cylinders I of $\mathbf{B}^\iota$, we have*

$$(6.1) \quad \kappa^{-1} < \frac{\mu_\iota(I)}{|I|_i^\delta e^{-nP}} < \kappa,$$

where i is a chart containing I of a bounded atlas $\mathcal{B}$ of $\mathcal{S}$;

- (ii) *We say that $\mathcal{S}$ is a C^{1+} realisation of a Gibbs measure $\nu = \nu_{\mathcal{S},\delta}$ if $\mu_\iota = (i_\iota)_* \nu_\iota$ where $\nu_\iota = (\pi_\iota)_* \nu$ and $\mu_\iota = \mu_{\mathcal{S},\delta}$ is a natural geometric measure of $\mathcal{S}$.*

Suppose that we have a C^{1+} self-renormalisable structure $\mathcal{S}$ on $\mathbf{B}^\iota$ and that $\mathcal{B}$ is a bounded atlas for it. Let $\delta > 0$. If I is a segment in $\mathbf{B}^\iota$, let $|I| = |I|_i$ be its length in any chart i of this atlas




FIGURE 13. The rectangles C_Λ^1, C_Λ^2 and $D_\Lambda^1, \dots, D_\Lambda^q$

which contains it. If C is a m -cylinder, let us denote m by $n(C)$ and $i_\ell(C)$ by I_C . For $m_1 \geq 1$ and $m_2 \geq 1$, let C be an m_1 -cylinder and D an m_2 -cylinder contained in the same 1-cylinder. Let

$$(6.2) \quad L_{\delta,s}(C : D) = \frac{\sum_{C' \subset C} |I_{C'}|^\delta e^{-n(C')s}}{\sum_{D' \subset D} |I_{D'}|^\delta e^{-n(D')s}}$$

where the sums are respectively over all cylinders contained in C and D and the values $|I_{C'}|$ and $|I_{D'}|$ are determined using the same chart in $\mathcal{B}$. Let the *pressure* $P = P(\mathcal{S}, \delta)$ be the infimum value of s for which the numerator (and the denominator) are finite.

If $\xi \in \Theta^{\ell'}$, then the leaf 1-cylinder segment $\xi_\Lambda = i(\pi_{\ell'}^{-1}\xi) \subset \Lambda$ is also regarded, without ambiguity, as a point in the train-track $\mathbf{B}^{\ell'}$. Similarly, if C is an n -cylinder of $\Theta^{\ell'}$ then the $(1, n)$ -rectangle $C_\Lambda = i(\pi_{\ell'}^{-1}C) \subset \Lambda$ is also regarded, without ambiguity, as an n -cylinder of the train-track $\mathbf{B}^{\ell'}$.

The following theorem follows from the results proved in [23]. It can also be deduced from standard results about Gibbs states such as those in [2].

Theorem 6.1. *Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on $\mathbf{B}^{\ell'}$. For every $\delta > 0$, there is a unique geometric natural measure $\mu_\ell = \mu_{\mathcal{S},\delta}$ with pressure $P = P(\mathcal{S}, \delta) \in \mathbb{R}$, and there is an unique τ -invariant Gibbs measure $\nu = \nu_{\mathcal{S},\delta}$ on Θ such that $\mu_\ell = (i_\ell)_*\nu_\ell$ where $\nu_\ell = (\pi_\ell)_*\nu$. Furthermore, the measure μ_ℓ has the following properties:*

- (i) *There is $0 < \alpha < 1$ such that if C and D are any two n -cylinders in $\Theta^{\ell'}$ such that I_C and I_D are contained in a common segment K then*

$$\frac{\mu_\ell(I_C)}{\mu_\ell(I_D)} \in (1 \pm \mathcal{O}(|K|^\alpha)) L_{\delta,P}(C : D) .$$

- (ii) *If $\rho : \text{msol}^{\ell'} \rightarrow \mathbb{R}$ is the extended measure scaling function of ν_ℓ , then*

$$\rho_\xi(C) = \lim_{m \rightarrow \infty} L_{\delta,P}(C_m : \xi_m) ,$$

where C_m and ξ_m are the cylinders given by $I_{C_m} = f_{\ell'}^m(C_\Lambda \cap \xi_\Lambda)$ and $I_{\xi_m} = f_{\ell'}^{m-1}\xi_\Lambda$.

- (iii) *(ratio decomposition) if C is an n -cylinder in $\Theta^{\ell'}$ and C_p is the primary cylinder containing C then*

$$(6.3) \quad \mu_\ell(I_C) = \int_{\xi \in \pi_{\ell'}(C)} \rho_\xi(C) \nu_{\ell'}(d\xi) .$$

Lemma 6.2. *Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on $\mathbf{B}^{\ell'}$ and let ρ be the extended measure scaling function of the Gibbs measure $\nu_{\mathcal{S},\delta}$.*

- (i) *If C and D are two cylinders contained in an n -cylinder E of $\Theta^{\ell'}$ then, for all ξ, η contained in the 1-cylinder $\pi_{\ell'}(\pi_{\ell'}^{-1}E)$ of $\Theta^{\ell'}$,*

$$(6.4) \quad \frac{\rho_\eta(C)}{\rho_\eta(D)} \in (1 \pm \mathcal{O}(\theta^n)) \frac{\rho_\xi(C)}{\rho_\xi(D)} .$$

(ii) Let $\mathbf{B}'$ be a no-gap train-track. Let $\xi, \eta \in \Theta'$ be such that the corresponding leaf segments in Λ have a common intersection K (or coincide). Let $(\xi : C^1), (\xi : C^2), (\eta : D^1), \dots, (\eta : D^q)$ be admissible pairwise distinct pairs in msc' such that (a) $\xi_\Lambda \cap C_\Lambda^1 = \xi_\Lambda \cap (\cup_{i=1}^p D_\Lambda^i) \subset K$, and (b) $\xi_\Lambda \cap C_\Lambda^2 = \xi_\Lambda \cap (\cup_{i=p+1}^q D_\Lambda^i) \subset K$ (see Figure 13). Then

$$(6.5) \quad \frac{\rho_\xi(C^1)}{\rho_\xi(C^2)} = \frac{\sum_{i=1}^p \rho_\eta(D^i)}{\sum_{i=p+1}^q \rho_\eta(D^i)}.$$

(iii) Let $\mathbf{B}^t$ be a no-gap train-track (and $\delta = 1$). Then for every admissible pair $(C : \xi) \in \text{msc}^t$ we get

$$(6.6) \quad \rho_\xi(C) = r^t(C_\Lambda \cap \xi_\Lambda : \xi_\Lambda)$$

where r^t is the ι -ratio function determined by the C^{1+} self-renormalisable structure.

Proof. *Proof of (i) and (ii).* Suppose that C and D are two cylinders contained in an n -cylinder E . Let E_1 be a $(n+1)$ -cylinder whose image under the shift map τ is E and let C_1 and D_1 be the cylinders in E_1 such that $\tau C_1 = C$ and $\tau D_1 = D$. Then

$$L_{\delta,P}(C_1 : D_1) \in (1 \pm \mathcal{O}(\theta^n)) L_{\delta,P}(C : D)$$

where (i) $0 < \theta < 1$ is independent of C, D, E and E_1 , and $P = P(\mathcal{S}, \delta)$ is the pressure. This follows directly from the definition of $L_{\delta,P}$ together with the fact that for all cylinders C', D' in E_1 ,

$$\frac{|I_{D'}|}{|I_{C'}|} \in (1 \pm \mathcal{O}(\theta^n)) \frac{|I_{\tau D'}|}{|I_{\tau C'}|}.$$

As a corollary of this we deduce (6.4). Then, equality (6.5) follows from using that the local holonomies are local diffeomorphisms in the self-renormalisable structure of $\mathbf{B}^t$.

Proof of (iii). In this case the self-renormalisable structure $\mathcal{S}$ is a local manifold structure as defined in Section 4.6 (i.e. the charts are homeomorphisms onto a subinterval of $\mathbb{R}$), and $\delta = 1$. Using (6.2), we get $P(\mathcal{S}, \delta) = 0$ and so the ratios $\mu(I)/|I|$ are uniformly bounded away from 0 and ∞ for all segments I in $\mathbf{B}^t$. Moreover, in this case, the length system ℓ matches in the sense that, if C is an n -cylinder then $\sum_{C'} |I_{C'}| = |I_C|$ where the sum is over all m -cylinders C' contained in C and $|I_C|$ and $|I_{C'}|$ are obtained using the same chart in $\mathcal{B}$. Thus, if C and D are n -cylinders and $I_C \cup I_D$ is a segment of $\mathbf{B}^t$ then

$$\frac{\mu_\iota(I_C)}{\mu_\iota(I_D)} \in (1 \pm \mathcal{O}(\theta^n)) \frac{|I_C|}{|I_D|}.$$

Hence,

$$\rho_\xi(C) = \lim_{m \rightarrow \infty} \frac{|f_{\iota'}^m(C_\Lambda \cap \xi_\Lambda)|}{|f_{\iota'}^m \xi_\Lambda|}$$

which implies (6.6).

Lemma 6.3. *If δ is the Hausdorff dimension of $\mathbf{B}^t$ then the ratios $\mu_\iota(I_C)/|I_C|^\delta$ are uniformly bounded away from 0 and ∞ . It follows from this that the Hausdorff δ -measure $\mathcal{H}^\delta$ is finite and positive on $\mathbf{B}^t$ and such that μ_ι is absolutely continuous with respect to $\mathcal{H}^\delta$.*

Proof. Since in this case δ is the Hausdorff dimension of $\mathbf{B}^t$, $P_\delta = 0$. Now, let us prove that the ratios $\mu_\iota(I)/|I|^\delta$ are uniformly bounded away from 0 and ∞ for all segments I . Suppose that I is any segment in $\mathbf{B}^t$. Then either there exists a cylinder C with $I_C \subset I$ such that $I \subset I_{mC}$ or there exist cylinders $I_C, I_D \subset I$ with a common endpoint such that $I \subset I_{mC} \cup I_{mD}$. In the first case

let $\hat{I} = I_C$ and $m\hat{I} = I_{mC}$, otherwise let $\hat{I} = I_C \cup I_D$ and $m\hat{I} = I_{mC} \cup I_{mD}$. Since by bounded geometry there exists $\sigma > 0$ such that for all cylinders $|I_C|/|I_{mC}| \in [\sigma, \sigma^{-1}]$,

$$\sigma|m\hat{I}| \leq |\hat{I}| \leq |I| \leq |m\hat{I}| \leq \sigma^{-1}|\hat{I}|.$$

By Theorem 6.1, the ratios $\mu_\iota(I_C)/|I_C|^\delta$ are uniformly bounded away from 0 and ∞ for cylinders C , and so the ratios $\mu_\iota(I)/|I|^\delta$ for segments I are also uniformly bounded away from 0 and ∞ .

Suppose that A is a subset of $\mathbf{B}^\iota$. Recall the definition of $\mathcal{H}_\varepsilon^\delta(A)$ as the infimum of the sums $\sum r_i^\delta$ where the r_i are the lengths of the segments of an ε -cover of A . Then the Hausdorff δ -dimensional outer measure of A is $\mathcal{H}^\delta(A) = \lim_{\varepsilon \rightarrow 0} \mathcal{H}_\varepsilon^\delta(A) = \sup_{\varepsilon > 0} \mathcal{H}_\varepsilon^\delta(A)$. Now suppose that A is a subset of $\mathbf{B}^\iota$ and $\{B_i\}$ is a cover of A by segments of length r_i . Then, $\mathcal{H}^\delta(A) \geq \mathcal{O}(\mu_\iota(A))$ because

$$\mathcal{H}^\delta(A) \geq \sum r_i^\delta \geq \mathcal{O}\left(\sum \mu_\iota(B_i)\right) \geq \mathcal{O}\left(\mu_\iota\left(\bigcup B_i\right)\right) \geq \mathcal{O}(\mu_\iota(A)).$$

Now suppose that A is a Borel subset of $\mathbf{B}^\iota$ with $\mu_\iota(A) > 0$. The set of segments in $\mathbf{B}^\iota$ is a Vitali class for A . Therefore, by the Vitali Covering Theorem (e.g. [4, 6]), given $\varepsilon > 0$, there is a countable disjoint sequence B_j of segments such that either $\sum_j |B_j|^\delta = \infty$ or $\mathcal{H}^\delta(A) \leq \sum_j |B_j|^\delta + \varepsilon$. But since the B_j 's are disjoint we get

$$\sum_j |B_j|^\delta \leq \mathcal{O}\left(\sum_j \mu_\iota(B_j)\right) \leq \mathcal{O}(\mu_\iota(A)).$$

Thus $\mathcal{H}^\delta(A) \leq \mathcal{O}(\mu_\iota(A)) + \varepsilon$. Letting $\varepsilon \rightarrow 0$ gives $\mathcal{H}^\delta(A) \leq \mathcal{O}(\mu_\iota(A))$. This proves that the measure μ_ι is proportional to $\mathcal{H}^\delta$. It follows immediately that $\mathcal{H}^\delta$ is positive and finite.

6.1. Measure ratio functions. In this subsection, we prove that, for every $\delta > 0$, a given C^{1+} self-renormalisable structure $\mathcal{S}$ on $\mathbf{B}^{\iota'}$ determines an ι -measure ratio function $\rho_{\mathcal{S},\delta}$ such that the Gibbs measure ν_ρ determined by $\rho_{\mathcal{S},\delta}$ (see Remark 5.8) is the same as the Gibbs measure $\nu_{\mathcal{S},\delta}$ which is C^{1+} realisable by the self-renormalisable structure $\mathcal{S}$.

Let $\xi \in \mathbf{B}^{\iota'}$ be an ι -leaf segment spanning of a Markov rectangle M . Let R be a rectangle inside M . There are cylinders $C_j \in \Theta^\iota$ such that $\pi_\iota R$ is the countable (or finite) union $\bigcup_{j \in \text{Ind}} I_{C_j}$ of cylinders $I_{C_j} = i_\iota(C_j)$ of $\mathbf{B}^\iota$, and any two of which intersect at most in a point of their boundary (see Figure 14). Suppose that $\xi \cap R \neq \emptyset$. Let $\xi' \in \Theta^{\iota'}$ be such that $i_\iota(\xi') = \xi$ (we note that ξ' might not be uniquely determined). Using (6.5), the following ratio is well-defined

$$(6.7) \quad \rho_{\iota,\xi}(R : M) = \sum_{j \in \text{Ind}} \rho_{\xi'}(C_j),$$

where $\rho_{\xi'}(C_j)$ is the ι -extended measure scaling function of $\nu_{\mathcal{S},\delta}$. If $\xi \cap R = \emptyset$ then we define $\rho_{\xi'}(R : M) = 0$. More generally, suppose that R_0 and R_1 are ι' -spanning rectangles contained in R . Then we define

$$\rho_{\iota,\xi}(R_0 : R_1) = \rho_{\iota,\xi}(R_0 : M) \rho_{\iota,\xi}(R_1 : M)^{-1}.$$

Theorem 6.4. (2-dimensional ratio decomposition) *Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure and $\mu_\iota = \mu_{\mathcal{S},\delta}$ a natural geometric measure for some $\delta > 0$. Suppose that R is a rectangle contained in a Markov rectangle M . Then*

$$(6.8) \quad \mu(R) = \int_{\pi_{\mathbf{B}^{\iota'}}(R)} \rho_{\iota,\xi}(R : M) \mu_{\iota'}(d\xi).$$

We now consider the case where $\mathbf{B}^\iota$ is a no-gap train-track. Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure and $\mu_\iota = \mu_{\mathcal{S},1}$ the natural measure (with pressure $P = 0$). Recall the definition of $t_R^{\iota'}$ as the set of spanning ι -leaf segments of the rectangle R (not necessarily a Markov rectangle). By the local




FIGURE 14. The holonomy injection t_θ .

product structure, one can identify t'_R with any spanning ι' -leaf segment $\ell^{\iota'}(x, R)$ of R . Suppose that R is a rectangle and M is a Markov rectangle and that $\theta : \ell = \ell^{\iota'}(x, R) \rightarrow \ell' \subset \ell^{\iota'}(x', M)$ is a basic holonomy defined on the spanning ι' -leaf segment ℓ . This defines an injection $t_\theta : t'_R \rightarrow t'_M$ which we call the *holonomy injection* induced by θ (see Figure 14). The measure $\mu_{\iota'}$ on $\mathbf{B}^{\iota'}$ induces a measure on t'_M which we can pull back to t'_R using t_θ to obtain a measure $\mu_{R,M}^\theta$ i.e. $\mu_{R,M}^\theta(E) = \mu_{\iota'}(\pi_{\mathbf{B}^{\iota'}}(t_\theta(E)))$.

Theorem 6.5. (2-dimensional ratio decomposition for SRB measures) *Let $\mathbf{B}^{\iota'}$ be a no-gap train-track. Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure and $\mu_\iota = \mu_{\mathcal{S},1}$ the natural measure (with pressure $P = 0$). If $t_\theta : t'_R \rightarrow t'_M$ is a holonomy injection as above with P a Markov rectangle then*

$$(6.9) \quad \mu(R) = \int_{t'_R} r_\iota(\xi : t_\theta(\xi)) \mu_{R,M}^\theta(d\xi) ,$$

where r_ι is the ι -ratio function determined by $\mathcal{S}$.

Remark 6.6. *Note that if $R \subset M$ then $t_\theta(\xi)$ is just the M -spanning ι -leaf containing ξ and $\mu_{R,M}^\theta = \mu_{\iota'}$.*

Since any rectangle can be written as the union of rectangles R with the property hypothesised in the theorem for some Markov rectangle, the above theorem gives an explicit formula for the measure of any rectangle in terms of a ratio decomposition using the ratio function which characterises the smooth structure of the train-track.

Proof of Theorems 6.4 and 6.5. Suppose that R is any rectangle, M is a Markov rectangle and $t_\theta : t'_R \rightarrow t'_M$ is a holonomy injection as above (in the case of Theorem 6.4 t_θ is the identity map). Then we note that there is $0 < \nu < 1$ such that for all $n > 0$ we can write $R = R_0 \cup \dots \cup R_{N(n)}$ where

- (i) $R_0, \dots, R_{N(n)}$ are rectangles which intersect at most in their boundary leaves and their spanning ι -leaf segments are also R -spanning ι -leaf segments;
- (ii) $P_i = t_\theta R_i$ and $\pi_{\iota'}(P_i)$ is an n -cylinder of $\mathbf{B}^{\iota'}$ for every $0 \leq i \leq N(n)$;

- (iii) R_0 is the empty set, or $\pi_{\iota'}(P_0)$ is strictly contained in an n -cylinder of $\mathbf{B}'$, and so, using the bounded geometry of the Markov map (see Section 4.3) and (6.1), $\mu(R_0) < \mathcal{O}(\varepsilon_0^n)$ for some $0 < \varepsilon_0 < 1$;

Let $S_i = f_{\iota'}^n R_i$ and $Q_i = f_{\iota'}^n P_i$ for $1 \leq i \leq N(n)$, and note that the rectangles Q_i are ι -spanning $(1, n)$ -rectangles of some Markov rectangle M_i . We note that if t_θ is not the identity there might be a non-empty set V_n of values of i such that S_i is not contained in the Markov rectangle M_i . However, since there are a finite number of Markov rectangles, the cardinality of the set V_n is bounded away from infinity, independently of $n \geq 0$. Hence, we disregard in what follows these values of $i \in V_n$, since the measure of the corresponding sets S_i converges to 0 when n tends to infinity. To prove the theorems we firstly note that by Lemma 6.2 and by (6.7) we obtain that, if Q_i, P_i and M_i are as above, for all $\xi, \eta \in M_i$,

$$\rho_\xi(S_i : Q_i) \in (1 \pm \mathcal{O}(\varepsilon^n)) \rho_\eta(S_i : Q_i),$$

for some $0 < \varepsilon < 1$. Thus, since $\mu(S_i) = \mu(\pi_\iota(S_i))$ and $\mu(Q_i) = \mu(\pi_\iota(Q_i))$ and by (6.3), if $\xi \in t_{M_i}'$,

$$\mu(S_i) \in (1 \pm \mathcal{O}(\varepsilon^n)) \rho_\xi(S_i : Q_i) \mu(Q_i).$$

Now consider the case of Theorem 6.4. Then, since R_i and P_i are contained in the same Markov rectangle, $\rho_\xi(S_i : Q_i)$ equals $\rho_{\xi_i}(R : M)$ for some $\xi_i \in t_{R_i}'$ and $\rho(Q_i) = \rho(P_i)$ which is equal to $\mu_{\iota'} P_i$ since P_i is an ι -spanning rectangle of the Markov rectangle M . Thus we have deduced that up to addition of a term that is $\mathcal{O}(\nu^n)$,

$$\mu(R) \in (1 \pm \mathcal{O}(\varepsilon^n)) \sum_{i=1}^{N(n)} \rho_{\xi_i}(R : M) \mu_{\iota'}(P_i).$$

Equation (6.8) follows on taking the limit $n \rightarrow \infty$.

Now consider the case of Theorem 6.5. Under its hypotheses we have that $\rho_\xi(S_i : Q_i) = r^\iota(\xi \cap S_i : \xi \cap Q_i)$ by (6.6) and (6.7). By the f -invariance of r^ι there is $\xi_i \in t_{R_i}'$ such that $r^\iota(\xi_i : t_\theta(\xi_i)) = r^\iota(\xi \cap S_i : \xi \cap Q_i)$. Thus, as above, we deduce that

$$\mu(R) \in (1 \pm \mathcal{O}(\varepsilon^n)) \sum_{i=1}^{N(n)} r^\iota(\xi_i : t_\theta(\xi_i)) \mu_{\iota'}(t_\theta R_i).$$

Equation (6.9) follows on taking the limit $n \rightarrow \infty$. $\square$

Lemma 6.7. *Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on $\mathbf{B}^\iota$ with natural measure $\mu_\iota = \mu_{\mathcal{S}, \delta}$ for some $\delta > 0$. Suppose that R is contained in a (n_s, n_u) -rectangle and that R' and R'' are ι' -spanning rectangles contained in R . Suppose in addition that either (i) R is contained in a Markov rectangle or (ii) $\mathbf{B}^\iota$ does not have gaps and there is a holonomy injection of R into a Markov rectangle (in this case $\delta = 1$ and $P = 0$). Then for every ι -leaf segment $\xi \in t_R'$ we have that*

$$(6.10) \quad \frac{\mu(R')}{\mu(R'')} \in (1 \pm \mathcal{O}(\varepsilon^{n_s + n_u})) \rho_\xi(R' : R'')$$

for some constant $0 < \varepsilon < 1$ independent of R, R', R'', n_s and n_u , (and in case (ii) $\rho_\xi(R : R') = r^\iota(\xi \cap R : \xi \cap R')$).

Proof. We give the proof for the second case since that for the first is similar. By Theorem 6.5, we have that

$$\frac{\mu(R)}{\mu(R')} = \frac{\int_{t_R'} r^\iota(\xi : t_\theta(\xi)) \mu_{R, M}^\theta(d\xi)}{\int_{t_{R'}} r^\iota(\xi : t_\theta(\xi)) \mu_{R', M}^\theta(d\xi)} = \frac{\int_{t_R'} r_\xi(R : R') r^\iota(\xi : t_\theta(\xi)) \mu_{R, M}^\theta(d\xi)}{\int_{t_{R'}} r^\iota(\xi : t_\theta(\xi)) \mu_{R', M}^\theta(d\xi)}.$$

where $r_\xi^\iota(R : R') = r^\iota(R \cap \xi : R' \cap \xi)$. Let $F = f_{\iota'}^{n_s+n_u}$. By inequality (3.2) (or inequality (6.4) in case (i)), there is $0 < \varepsilon < 1$ such that

$$r_{F\eta}^\iota(FR : FR') \in (1 \pm \mathcal{O}(\varepsilon^{n_s+n_u}))r_{F\xi}^\iota(FR : FR')$$

for all $\xi, \eta \in t_R^{\iota'}$. Thus,

$$r_\eta^\iota(R : R') \in (1 \pm \mathcal{O}(\varepsilon^{n_s+n_u}))r_\xi^\iota(R : R')$$

and so

$$\frac{\mu(R)}{\mu(R')} \in (1 \pm \mathcal{O}(\varepsilon^{n_s+n_u}))r_\xi^\iota(R : R') .$$

Similarly,

$$\frac{\mu(R)}{\mu(R'')} \in (1 \pm \mathcal{O}(\varepsilon^{n_s+n_u}))r_\xi^\iota(R : R') .$$

Putting together the previous two equations we obtain (6.10).

Lemma 6.8. *Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on $\mathbf{B}^\iota$ with natural measure $\mu_\iota = \mu_{\mathcal{S},\delta}$ for some $\delta > 0$. The values*

$$\rho_{\mathcal{S},\delta}(\xi \cap R' : \xi \cap R'') = \rho_\xi(R' : R'')$$

(as in Lemma 6.7) determine an ι -measure ratio function $\rho_{\mathcal{S},\delta}$ with the following properties: (i) The Gibbs measure ν_ρ determined by the ι -measure ratio function $\rho_{\mathcal{S},\delta}$ (see Remark 5.8) is the same as the Gibbs measure $\nu_{\mathcal{S},\delta}$ which is C^{1+} realisable by the self-renormalisable structure $\mathcal{S}$; (ii) If $\mathbf{B}^\iota$ is a no-gap train-track then $\rho_{\mathcal{S},1} = r$, where r is the ratio function determined by the C^{1+} self-renormalisable structure $\mathcal{S}$.

Proof. Let us prove this lemma first in the case where the train-track $\mathbf{B}^\iota$ does not have gaps and then in the case where the train-track $\mathbf{B}^\iota$ has gaps.

(i) $\mathbf{B}^\iota$ does not have gaps. Then $\delta = 1$ and, by Lemma 6.2 (iii), we have $\rho_\xi(R' : R'') = r(\xi \cap R' : \xi \cap R'')$ where r is the ratio function determined by the C^{1+} self-renormalisable structure $\mathcal{S}$. Hence $\rho_{\mathcal{S},\delta} = r$ is an ι -measure ratio function. Using (6.10), we get that the Gibbs measure, which is a C^{1+} realisation of the natural geometric measure $\mu_{\mathcal{S},\delta}$, determines an ι -measure solenoid function which induces the ι -measure ratio function $\rho_{\mathcal{S},\delta}$.

(ii) $\mathbf{B}^\iota$ has gaps. Let $\xi, \eta \in \mathbf{B}^{\iota'}$ be boundaries of Markov rectangles such that $\xi \cap \eta \neq \emptyset$. Let M, M', R and R' be rectangles such that $\xi \cap R = \eta \cap M$ and $\xi \cap R' = \eta \cap M'$. Using (6.5) and (6.7), we get that

$$\rho_\xi(R : R') = \rho_\eta(M : M') .$$

Therefore, the ι -measure ratio function $\rho_{\mathcal{S},\delta}$ is well-defined for every ι -leaf segments I and J contained in a spanning leaf segment of a Markov rectangle. By f -invariance of μ and (6.10), we get that

$$(6.11) \quad \rho_\xi(R : R') = \rho_{f_{\iota'}\xi}(f_{\iota'}R : f_{\iota'}R')$$

is invariant under f . Let I and J be ι -leaf segments such that (a) there is an ι -leaf segment K such that $I, J \subset K$, and (b) I or J has non-empty interior. Then there is $n > 0$, $\xi \in \mathbf{B}^{\iota'}$, R and R' such that $f_{\iota'}^n I = \xi \cap R$ and $f_{\iota'}^n J = \xi \cap R'$. Hence, using (6.11), the ratio

$$\rho_{\mathcal{S},\delta}(I : J) = \rho_\xi(R : R')$$

is well defined independently of n . Using (6.10), we get that (5.1) is satisfied and the Gibbs measure, which is a C^{1+} realisation of the natural geometric measure $\mu_{\mathcal{S},\delta}$, determines an ι -measure solenoid function which induces the ι -measure ratio function $\rho_{\mathcal{S},\delta}$.




FIGURE 15. The rectangles $A_m = [I, a_m]$ and $B_m = [K, b_m]$.

6.2. Dual measure ratio function. We will show that an ι -measure ratio function ρ_ι determines an unique dual function $\rho_{\iota'}$ which is an ι' -measure ratio function.

Definition 6.2. We say that the ι -measure ratio function ρ_ι and the ι' -measure ratio function $\rho_{\iota'}$ are dual if both determine the same Gibbs measure $\nu = \nu_{\rho_\iota} = \nu_{\rho_{\iota'}}$ on Θ (see Remark 5.8).

Lemma 6.9. Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on $\mathbf{B}^\iota$ with ι -measure ratio function $\rho_\iota = \rho_{\mathcal{S},\delta}$ corresponding to the Gibbs measure $\nu = \nu_{\mathcal{S},\delta}$. Then there is an unique ι' -measure ratio function $\rho_{\iota'}$ determining the same Gibbs measure $\nu_{\rho_{\iota'}} = \nu$ on Θ .

Proof. Let $\mu = i_*\nu$. The dual $\rho_{\iota'}$ of ρ_ι is constructed as follows: Let I and K be (i) two ι' -leaf segments contained in a common n -cylinder ι' -leaf, or also (ii) two ι' -leaf segments contained in a union of two n -cylinders with a common endpoint in the case of a local manifold structure. Choose $p \in I$ and $p' \in K$. Let a_m be the ι -leaf N -cylinders containing p , and b_m the ι -leaf containing p' and holonomic to a_m . Let $A_m = [I, a_m]$ and $B_m = [K, b_m]$ (see Figure 15). Now, let us prove that

(i)

$$(6.12) \quad \frac{\mu(A_{m+1})}{\mu(B_{m+1})} \in (1 \pm \mathcal{O}(\varepsilon^{n+m})) \frac{\mu(A_m)}{\mu(B_m)}$$

for some $0 < \varepsilon < 1$;

(ii) the dual measure ratio function is given by

$$(6.13) \quad \rho_{\iota'}(I : K) = \lim_{m \rightarrow \infty} \frac{\mu(A_m)}{\mu(B_m)};$$

By Lemma 6.7, there is $0 < \varepsilon < 1$ such that

$$\mu(A_{m+1})/\mu(A_m) \in (1 \pm \mathcal{O}(\varepsilon^{n+m}))\rho_\iota(a_{m+1} : a_m),$$

and, similarly,

$$\mu(B_{m+1})/\mu(B_m) \in (1 \pm \mathcal{O}(\varepsilon^{n+m}))\rho_\iota(b_{m+1} : b_m).$$

Since ρ_ι is an ι -ratio function,

$$\rho_\iota(a_{m+1} : a_m) \in \mathcal{O}(\varepsilon^{n+m})\rho_\iota(b_{m+1} : b_m).$$

Therefore, (6.12) follows. Furthermore, (6.12) implies (6.13).

Using (6.12), we obtain that $\rho_{\iota'}$ is an ι' -measure ratio function: $\rho_{\iota'}$ is f -invariant, $\rho_{\iota'}(I : K) = \rho_{\iota'}(K : I)^{-1}$ and

$$\rho_{\iota'}(I : K) = \rho_{\iota'}(I_1 : K) + \rho_{\iota'}(I_2 : K)$$

for ι -leaf segments I_1 and I_2 with at most one common point and such that $I = I_1 \cup I_2$. Again using (6.12), $\rho_{\iota'}$ satisfies inequality (5.1).

7. COCYCLE-GAP PAIRS

In this section, we are going to introduce the gap-cocycle pairs (γ, J) consisting of a gap ratio function γ and of a measure-length ratio cocycle J satisfying a gap-cocycle property. In Section 8, we will apply this description to give an explicit geometric construction of all C^{1+} self-renormalisable structures and all C^{1+} hyperbolic diffeomorphisms which have natural geometric measures.

7.1. Measure-length ratio cocycle. Let $\mathbf{B}^\iota$ be a gap train-track. For each Markov rectangle R let t_R^ι be the set of ι -segments of R . Let us denote by $\mathbf{B}_o^{\iota'}$ the disjoint union $\sqcup_{i=1}^m t_{R_i}^{\iota'}$ over all Markov rectangles $R_1, \dots, R_m$ (without doing any extra-identification). In this section, for every $\xi \in \mathbf{B}_o^{\iota'}$ and $n \geq 1$, we denote by ξ_n the n -cylinder $\pi_\iota f_\iota^{n-1} \xi$ of $\mathbf{B}^\iota$.

Definition 7.1. Let $\mathbf{B}^\iota$ be a gap train-track and ρ be a ι -measure ratio function. We say that $J : \mathbf{B}_o^{\iota'} \rightarrow \mathbb{R}^+$ is a (ρ, δ, P) ι -measure-length ratio cocycle if $J = \kappa / (\kappa \circ f_\iota)$ where κ is a positive Hölder continuous function on $\mathbf{B}_o^{\iota'}$ and is bounded away from 0, and

$$(7.1) \quad \sum_{f_\iota \eta = \xi} J(\eta) \rho(f_\iota \eta : m(f_\iota \eta))^{1/\delta} e^{P/\delta} < 1,$$

for every $\eta \in \mathbf{B}_o^{\iota'}$.

We note that in (7.1), the mother of η is not defined because η is a leaf primary cylinder segment, and so we used instead the mother of the leaf 2-cylinder $f_\iota \eta$.

Let us consider a C^{1+} self-renormalisable structure $\mathcal{S}$ on $\mathbf{B}^\iota$, and fix a bounded atlas $\mathcal{B}$ for $\mathcal{S}$. Let $\delta > 0$. By Theorem 6.1, the C^{1+} self-renormalisable structure $\mathcal{S}$ C^{1+} -realises a Gibbs measure $\nu = \nu_{\mathcal{S}, \delta}$ as a natural invariant measure $\mu = \mu_{\mathcal{S}, \delta} = i_* \nu$ with pressure $P = P(\mathcal{S}, \delta)$. Let $\rho = \rho_{\mathcal{S}, \delta}$ be the corresponding ι -measure ratio function (see Lemma 6.8). Since μ is a natural geometric measure, for every $\xi \in \mathbf{B}_o^{\iota'}$, the ratios $|\xi_n|_i e^{-nP/\delta} / \mu(\xi_n)^{1/\delta}$ are uniformly bounded away from 0 and ∞ , where the length $|\xi_n|_i$ is measured in any chart $i \in \mathcal{B}$ containing ξ_n in its domain. Therefore,

$$\kappa_i(\xi_n) = \frac{|\xi_n|_i e^{-nP/\delta}}{\mu(\xi_n)^{1/\delta}}$$

is well-defined. By Lemma 6.7, we get

$$\frac{\mu_\iota(\xi_n)}{\mu_\iota(m\xi_n)} \in (1 \pm \mathcal{O}(\varepsilon^n)) \rho(f_\iota \xi : m(f_\iota \xi))$$

for some $0 < \varepsilon < 1$. Hence, the ratios $\mu_\iota(\xi_n) / \mu_\iota(m\xi_n)$ converge exponentially fast along backward orbits ξ of cylinders. By (4.1), we get that $|\xi_n| / |m\xi_n|$ also converge exponentially fast along backward orbits ξ of cylinders. Therefore, it follows that there is a Hölder function $J_{\mathcal{S}, \delta} : \mathbf{B}_o^{\iota'} \rightarrow \mathbb{R}$ such that

$$(7.2) \quad \frac{\kappa_i(\xi_n)}{\kappa_i(m\xi_n)} \in (1 \pm \mathcal{O}(\varepsilon^n)) J_{\mathcal{S}, \delta}(\xi)$$

for some $0 < \varepsilon < 1$.

Lemma 7.1. Let $\mathbf{B}^\iota$ be a gap train-track. Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure, and $\delta > 0$. Let $\mu_{\mathcal{S}, \delta}$ be the natural geometric measure with pressure $P = P(\mathcal{S}, \delta)$, and $\rho = \rho_{\mathcal{S}, \delta}$ the corresponding ι -measure ratio function. The function $J_{\mathcal{S}, \delta} : \mathbf{B}_o^{\iota'} \rightarrow \mathbb{R}^+$ given by (7.2) is a (ρ, δ, P) ι -measure-length ratio cocycle.

Proof. If I is an n -cylinder in $\mathbf{B}^l$, then $\sum_{mI'=I} |I'| < |I|$, where the lengths are measured in the same chart. Thus, since $|I'| = \kappa_i(I') \mu_l(I')^{1/\delta} e^{(n+1)P/\delta}$ we deduce that

$$\sum_{mI'=I} \frac{\kappa_i(I')}{\kappa_i(I)} \left(\frac{\mu_l(I')}{\mu_l(I)} \right)^{1/\delta} e^{P/\delta} < 1.$$

For every $\xi \in \mathbf{B}'_o$, we have that $\tau_{l'}\eta = \xi$ if, and only if, $\eta_{n+1} \subset \xi_n$ for every $n \geq 1$. Hence, the Hölder continuous function $J = J_{\mathcal{S},\delta}$ satisfies (7.1).

Now, suppose that $\xi \in \mathbf{B}'_o$ is such that there exists $p \geq 1$ with the property that $\xi_{np} \subset \xi_n$ for every $n \geq 1$. By (7.2), we get

$$\begin{aligned} \frac{\kappa_{i_0}(\xi_{jp})}{\kappa_{i_0}(\xi_{(j-1)p})} &= \prod_{l=0}^{p-1} \frac{\kappa_{i_{l+1}}(\xi_{(j-1)p+l+1})}{\kappa_{i_l}(\xi_{(j-1)p+l})} \\ &\in (1 \pm \mathcal{O}(\nu^{(j-1)p})) \prod_{l=0}^{p-1} J(f_{l'}^l(\xi)) \end{aligned}$$

where $i_0, \dots, i_{p-1}$ are charts contained in a bounded atlas of $\mathcal{S}$. Thus, for all $1 < m < M$, we have

$$\begin{aligned} \frac{\kappa_{i_0}(\xi_{Mp})}{\kappa_{i_0}(\xi_{mp})} &= \prod_{n=0}^{M-m-1} \frac{\kappa_{i_0}(\xi_{(n+m+1)p})}{\kappa_{i_0}(\xi_{(n+m)p})} \\ &\in (1 \pm \mathcal{O}(\nu^{mp})) \left[\prod_{l=0}^{p-1} J(f_{l'}^l(\xi)) \right]^{M-m}. \end{aligned}$$

Since the term on the left of this equation is uniformly bounded away from 0 and ∞ , it follows that $\prod_{l=0}^{p-1} J(f_{l'}^l(\xi)) = 1$. From Livšic's theorem (e.g. see [12]) we get that $J_{\mathcal{S},\delta} = \kappa/(\kappa \circ f_{l'})$ where κ is a positive Hölder continuous function on $\mathbf{B}'_o$ and is bounded away from 0.

7.2. Gap ratio function. Let $\mathbf{B}^l$ be a gap train-track. Let $\mathcal{G}'$ be the set of all pairs $(\xi_1 : \xi_2) \in \mathbf{B}'_o \times \mathbf{B}'_o$ such that $mf_{l'}\xi_1 = mf_{l'}\xi_2$. The metric d_Λ induces a natural metric $d_{\mathcal{G}'}$ on $\mathcal{G}'$ given by

$$d_{\mathcal{G}'}((\xi_1 : \xi_2), (\eta_1 : \eta_2)) = \max\{d_\Lambda(\xi_1, \eta_1), d_\Lambda(\xi_2, \eta_2)\}.$$

Definition 7.2. A function $\gamma : \mathcal{G}' \rightarrow \mathbb{R}^+$ is an ι -gap ratio function if it satisfies the following conditions:

- (i) $\gamma(\xi_1 : \xi_2)$ is uniformly bounded away from 0 and ∞ ;
- (ii) $\gamma(\xi_1 : \xi_2) = \gamma(\xi_1 : \xi_3)\gamma(\xi_3 : \xi_2)$;
- (iii) there are $0 < \theta < 1$ and $C > 1$ such that

$$(7.3) \quad |\gamma(\xi_1 : \xi_2) - \gamma(\eta_1 : \eta_2)| \leq C (d_{\mathcal{G}'}((\xi_1 : \xi_2), \gamma(\eta_1 : \eta_2)))^\theta.$$

We note that part (ii) of this definition implies that $\gamma(\xi_1 : \xi_2) = \gamma(\xi_2 : \xi_1)^{-1}$.

Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on $\mathbf{B}^l$ and $\mathcal{B}$ a bounded atlas for $\mathcal{S}$. Then the gap ratio function $\gamma_{\mathcal{S}}$ is well-defined by

$$(7.4) \quad \gamma_{\mathcal{S}}(\xi : \eta) = \lim_{n \rightarrow \infty} \frac{|\pi_{\mathbf{B}^l} f_l^n \xi|_{i_n}}{|\pi_{\mathbf{B}^l} f_l^n \eta|_{i_n}}$$

where $i_n \in \mathcal{B}$ contains in its domain the n -cylinder $mf_l^n \xi$ (we note that $mf_l^n \xi = mf_l^n \eta$).

7.3. Ratio functions. We are going to construct the ratio function of a C^{1+} self-renormalisable structure from the gap ratio function and measure-length ratio cocycle.

Lemma 7.2. *Let $\mathbf{B}^\iota$ be a gap train-track. Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure. Let $r_{\mathcal{S}}$ be the corresponding ι -ratio function. Let $\delta > 0$ and let $\mu = \mu_{\mathcal{S},\delta}$ be the natural geometric measure with pressure $P = P(\mathcal{S},\delta)$ and $\rho_{\mathcal{S},\delta}$ the corresponding ι -measure ratio function. Let $J_{\mathcal{S},\delta}$ and $\gamma_{\mathcal{S}}$ be the corresponding ι -gap ratio function and ι -measure-length ratio cocycle. Then the following equalities are satisfied:*

(i) *Let I be an ι -leaf n -cylinder contained in the ι -leaf $(n-1)$ -cylinder L . Then*

$$(7.5) \quad r_{\mathcal{S}}(I : L) = J_{\mathcal{S},\delta}(\xi_I) \rho_{\mathcal{S},\delta}(I : L)^{1/\delta} e^{P/\delta}$$

where $\xi_I = f_\iota^{n-1}I \in \mathbf{B}_o^\iota$.

(ii) *Let I be an n -cylinder and K an n -gap and both contained in a $(n-1)$ -cylinder L . Then*

$$(7.6) \quad r_{\mathcal{S}}(I : K) = r_{\mathcal{S}}(I : L) \frac{\sum_{G \subset L} \gamma_{\mathcal{S}}(G : K)}{1 - \sum_{D \subset L} r_{\mathcal{S}}(D : L)}.$$

where the sum in the numerator is over all n -gaps $G \subset L$ and the sum in the denominator is over all n -cylinders $D \subset L$.

Proof. For every n -cylinder $I \subset \mathbf{B}^\iota$, define $\kappa_i(I) = |I|_i e^{-nP/\delta} / \mu_\iota(I)^{1/\delta}$ and let $J_{\mathcal{S},\delta}$ be the associate measure-length cocycle. Let I be an ι -leaf n -cylinder, L the ι -leaf $(n-1)$ -cylinder containing I . Choose $p \in I$ and let U_m be the ι' -leaf m -cylinders containing p . Let A_m be the rectangle $[I, U_m]$ and B_m be the rectangle $[L, U_m]$. Then $f_\iota^{m-1}A_m$ and $f_\iota^{m-1}B_m$ are ι' -spanning rectangles of some Markov rectangle. Let a_m and b_m be the projections of these into $\mathbf{B}^\iota$. Then by the invariance of μ , $\mu(A_m)/\mu(B_m) = \mu_\iota(a_m)/\mu_\iota(b_m)$ and therefore

$$\begin{aligned} \rho_{\mathcal{S},\delta}(I : L)^{1/\delta} &= \lim_{m \rightarrow \infty} \frac{\mu(A_m)^{1/\delta}}{\mu(B_m)^{1/\delta}} \\ &= \lim_{m \rightarrow \infty} \frac{\mu_\iota(a_m)^{1/\delta}}{\mu_\iota(b_m)^{1/\delta}} \\ &= \lim_{m \rightarrow \infty} \frac{\kappa(a_m)^{-1} |a_m|_{i_m} e^{-(n+m)P/\delta}}{\kappa(b_m)^{-1} |b_m|_{i_m} e^{-(n+m-1)P/\delta}} \\ &= J_{\mathcal{S},\delta}(\xi_I)^{-1} r_{\mathcal{S}}(I : L) e^{-P/\delta} \end{aligned}$$

where $|a_m|_{i_m}$ and $|b_m|_{i_m}$ are measured in a chart i_m of the bounded atlas on $\mathbf{B}^\iota$, and ξ_I is the leaf primary cylinder segment $f_\iota^{n-1}(I)$. Thus, equation (7.5) is satisfied.

We note that the ratio of the size of K to the size ℓ of the totality of gaps G in L is given by $(\sum_{G \subset L} \gamma_{\mathcal{S}}(G : K))^{-1}$ where $\gamma_{\mathcal{S}}$ is the gap ratio function and the sum is over all n -gaps in L . But since the complement of the gaps in L is the union of n -cylinders we have that the ratio of ℓ to the size of L is $1 - \sum_{D \subset L} r_{\mathcal{S}}(D : L)$ where the sum is over all n -cylinders D in L . Thus we deduce that for $r_{\mathcal{S}}(I : K)$ we should take

$$(7.7) \quad r_{\mathcal{S}}(I : K) = r_{\mathcal{S}}(I : L) \frac{\sum_{G \subset L} \gamma_{\mathcal{S}}(G : K)}{1 - \sum_{D \subset L} r_{\mathcal{S}}(D : L)}$$

which proves (7.6).

7.4. Cocycle-gap pairs. In this section, we are going to construct a *gap-cocycle* map b which reflects the cylinder-gap condition of an ι -solenoid function (see Section 3.9), i.e the ratios are well-defined along the ι -boundaries of the Markov rectangles. Hence, r is an ι -ratio function.

Let $\mathbf{B}^\iota$ be a gap train-track and $\mathbf{B}'$ a no-gap train-track (as in the case of codimension one attractors or repellers). Let $\mathcal{Q}$ be the set of all periodic orbits O which are contained in the ι -boundaries of the Markov rectangles. For every periodic orbit $O \in \mathcal{Q}$, let us choose a point $x = x(O)$ belonging to the orbit O . Let us denote by $p(x)$ the smallest period of x . Let us denote by $M(1, x)$ and $M(2, x)$ the Markov rectangles containing the point x . Let us denote by $l_i(x)$ the ι -leaf i -cylinder segment of Markov rectangle $M(1, x)$ containing the point x . Let $A(f_\iota^i(x))$ be the smallest ι -leaf segment containing all the ι -boundary leaf segments of Markov rectangles intersecting the global leaf segment passing through the point $f_\iota^i(x)$. Let $q(x)$ be the smallest integer which is a multiple of $p(x)$, such that

$$A(f_\iota^i(x)) \subset f_\iota^{q(x)+i}(l_i(x))$$

for every $0 \leq i < p(x)$. Let us denote the ι -leaf segments $f_\iota^{q(x)+i}(l_i(x))$ by $L_i(x)$. We note that when using the notation $L_i(x)$, we will always consider i to be $i \bmod p(x)$. For every $j \in \{1, 2\}$, let $J(j, x)$ be the primary ι' -leaf segment contained in $M(j, x)$ with x as an endpoint such that $R(j, i, x) = [f_\iota^{q(x)+i}(l_i(x)), f_\iota^{q(x)+i}(J(j, x))]$ is a rectangle for every $0 \leq i < p(x)$. Let $\text{Co}(j, i, x) \subset \mathbf{B}'_o$ be the set of all ι -primary leaves ξ of Markov rectangles M such that $f_\iota \xi \subset L_i(x)$ and $f_\iota M \cap R(j, i, x)$ has non-empty interior. Let $\text{Gap}(j, i, x) \subset \mathcal{G}'$ be the set of all sister pairs (ξ_1, ξ_2) such that $m f_\iota \xi_1 (= m f_\iota \xi_2)$ is an ι -primary leaf of a Markov rectangle M with the property that $M \cap R(j, i, x)$ has non-empty interior. Let $\text{Co}_j = \cup_{O \in \mathcal{Q}} \cup_{i=0}^{p(x(O))-1} \text{Co}(j, i, x(O))$ and $\text{Gap}_j = \cup_{O \in \mathcal{Q}} \cup_{i=0}^{p(x(O))-1} \text{Gap}(j, i, x(O))$. Let ρ be an ι -measure ratio function with corresponding Gibbs measure ν . Let $\mathcal{D}_j(\rho, \delta, P)$ be the set of all pairs (γ_j, J_j) with the following properties:

- (i) $\gamma_j : \text{Gap}_j \rightarrow \mathbb{R}^+$ is a map;
- (ii) $J_j : \text{Co}_j \rightarrow \mathbb{R}^+$ is a map satisfying property (7.1), with respect to (ρ, δ, P) , for every $\xi \in \text{Co}_j$ such that $\xi \subset \cup_{O \in \mathcal{Q}} \cup_{i=0}^{p(x(O))-1} L_i(x(O))$.
- (iii) For every $x(O) \in \mathcal{Q}$, letting $x = x(O)$, $\prod_{l=0}^{p(x)-1} J_j(f_\iota^l I^j(i, x)) = 1$, where $I^j(i, x) \subset \text{Co}_j$ is a ι' -primary leaf segment containing the periodic point $f_\iota^i(x)$.

For every $x(O) \in \mathcal{Q}$, let $x = x(O)$ and let $A(1, x)$ and $A(2, x)$ be the $p(x)$ -cylinders of $M(1, x)$ and $M(2, x)$, respectively, containing the point x . The points

$$\pi_{\mathbf{B}'_o} A(j, x), \pi_{\mathbf{B}'_o} f_\iota A(j, x), \dots, \pi_{\mathbf{B}'_o} f_\iota^{p(x)-1} A(j, x)$$

in $\mathbf{B}'_o$ form a periodic orbit, under f_ι , with period $p(x)$, where $\pi_{\mathbf{B}'_o} : \Lambda \rightarrow \mathbf{B}'_o$ is the natural projection. The primary cylinders contained in the sets $\text{Co}(j, i, x)$ are pre-orbits of the points $\pi_{\mathbf{B}'_o} f_\iota^i A(j, x)$ in $\mathbf{B}'_o$, under f_ι . Hence, we note that, if $\prod_{i=0}^{p(x)-1} J(\pi_{\mathbf{B}'_o} f_\iota^i A(j, x)) = 1$, then, by Livšic's theorem (e.g. see [12]), there is a map k such that, for every $\xi \in \text{Co}(j, i, x)$, $J(\xi) = k(\xi)/(k \circ f_\iota)(\xi)$.

We say that C is an *out-gap segment of a rectangle* R if C is a gap segment of R and is not a leaf n -gap segment of any Markov rectangle M such that $M \cap R$ is a rectangle with non-empty interior.

We say that C is a *leaf n -cylinder segment of a rectangle* R , if C is a leaf n -cylinder segment of a Markov rectangle M such that $M \cap R$ is a rectangle with non-empty interior. We say that C is a *leaf n -gap segment of a rectangle* R , if C is a leaf n -gap segment of a Markov rectangle M such that $M \cap R$ is a rectangle with non-empty interior. We say that C is an *n -leaf segment of a rectangle* R , if C is a leaf n -cylinder segment of R or if C is a leaf n -gap segment of R .

Lemma 7.3. *Let $(\gamma_1, J_1) \in \mathcal{D}_1(\rho, \delta, P)$. Let $x = x(O)$, where $O \in \mathcal{Q}$. For every $i \in \{0, 1, \dots, p(x) - 1\}$ and for all 2-leaf segments $C \subset L_i(x)$ of $R(1, i, x)$, the ratios $r(C : mC)$ are uniquely determined*

such that they are invariant under f , satisfy the matching condition, and satisfy equalities (7.5) and (7.6).

Proof. If $C \subset L_i(x)$ is a leaf 2-cylinder segment of $R(1, i, x)$, then we define the ratio $r(C : mC)$, using (7.5), by

$$(7.8) \quad r(C : mC) = J(\xi_C) \rho(C : mC)^{1/\delta} e^{P/\delta}$$

where $\xi_C = f_i C \in \text{Co}_1$. For every sister pair $(\xi_1 : \xi_2) \in \text{Gap}_1$ we define the ratio $r(f_{i'} \xi_1 : f_{i'} \xi_2)$ equal to $\gamma(\xi_1 : \xi_2)$. If $C \subset L_i(x)$ is a leaf 2-gap segment of $R(1, i, x)$, then we define the ratio $r(C : mC)$ by

$$(7.9) \quad r(C : mC) = \frac{1 - \sum_{D \subset mC} r(D : mC)}{\sum_{G \subset mC} r(G : C)},$$

where the sum, in the numerator, is over all 2-cylinders $D \subset mC$ of $R(1, i, x)$, and the sum, in the denominator, is over all 2-gaps $G \subset mC$ of $R(1, i, x)$. Hence,

$$\sum_{C \subset mC} r(C : mC) = 1$$

where the sum is over all 2-leaf segments $C \subset mC$ of $R(1, i, x)$.

Lemma 7.4. *Let $(\gamma_1, J_1) \in \mathcal{D}_1(\rho, \delta, P)$. Let $x = x(O)$, where $O \in \mathcal{Q}$, and let $i \in \{0, 1, \dots, p(x)-1\}$. For all $n \geq 0$, and for all out-gaps and all 2-leaf segments $C \subset f_i^{n+i} \ell_i(x)$ of $f_i^{n+i} M(1, x)$, the ratios $r(C : f_i^{n+i} \ell_i(x))$ are uniquely determined such that they are invariant under f , satisfy the matching condition, and satisfy equalities (7.5) and (7.6).*

Proof. Let us denote $f_i^n M(1, x)$ by M_n and $f_i^n \ell_i(x)$ by L_i^n . The proof follows by induction on $n \geq 0$. For the case $n = 0$, the ratios $r(C : L_i^{n+i})$ are uniquely determined by Lemma 7.3. Let us prove that the ratios $r(C : L_i^{n+1+i})$ are uniquely determined using the induction hypotheses with respect to n . For every out-gap and every primary cylinder segment $C \subset L_i^{n+1+i}$ of $f_i^{n+i} M(1, x)$, $f_{i'} C$ is a out-gap or a 2-leaf segment. Hence, by the induction hypotheses, the ratio $r(f_{i'} C : L_i^{n+i})$ is well-defined. Therefore, using the invariance of f , we define

$$(7.10) \quad r(C : L_i^{n+1+i}) = r(f_{i'} C : L_i^{n+i}).$$

For every 2-leaf segment $C \subset L_i^{n+1+i}$ of $f_i^{n+i} M(1, x)$, the ratio $r(C : mC)$ is well-defined by Lemma 7.3. Hence, by (7.10), we define

$$r(C : L_i^{n+1+i}) = r(C : mC) r(mC : L_i^{n+1+i})$$

which ends the proof of the induction.

Lemma 7.5. *Let $(\gamma_1, J_1) \in \mathcal{D}_1(\rho, \delta, P)$. Let $x = x(O)$, where $O \in \mathcal{Q}$, and let $i \in \{0, 1, \dots, p(x)-1\}$. Let $n \geq 0$ and $j \in \{0, \dots, n\}$. For all out-gaps and all $j+2$ -leaf segments $C \subset f_{i'}^n L_i(x)$ of $f_{i'}^n R(1, i, x)$, the ratios $r(C : f_{i'}^n L_i(x))$ are uniquely determined such that they are invariant under f , satisfy the matching condition, and satisfy equalities (7.5) and (7.6).*

Proof. The proof follows by induction in $n \geq 0$. For the case $n = 0$, noting that $L_i(x) = f_i^{q(x)+i} \ell_i(x)$, the ratios $r(C : L_i(x))$ are well-defined by Lemma 7.4. Hence, using the matching condition, the ratio $r(f_{i'}^{n+1} L_{i+1}(x) : f_{i'}^n L_i(x))$ is well-defined. Let us prove that for all out-gaps and $j+2$ -leaf segments $C \subset f_{i'}^{n+1} L_i(x)$ of $f_{i'}^{n+1} R(1, i, x)$, with $1 \leq j \leq n+1$, the ratios $r(C : f_{i'}^{n+1} L_i(x))$ are uniquely determined using the induction hypotheses with respect to n . By the induction hypotheses and by the matching condition, the ratio $r(f_i C : f_{i'}^n L_i(x))$ is well-defined. By invariance of f , we define $r(C : f_{i'}^{n+1} L_i(x)) = r(f_i C : f_{i'}^n L_i(x))$. which ends the proof of the induction.

Let us attribute the ratios for the cylinders and gaps of $R(2, i, x)$ such that they agree with the ratios previously defined in $R(1, i, x)$.

Lemma 7.6. *Let $(\gamma_1, J_1) \in \mathcal{D}_1(\rho, \delta, P)$. Let $x = x(O)$, where $O \in \mathcal{Q}$, and let $i \in \{0, 1, \dots, p(x)-1\}$. Let $n \geq 0$ and $j \in \{1, \dots, n\}$. For all out-gaps and all $j+2$ -leaf segments $C \subset f_{\iota'}^n L_i(x) \setminus f_{\iota'}^{n+1} L_{i+1}(x)$ of $f_{\iota'}^n R(2, i, x)$, the ratios $r(C : f_{\iota'}^n L_i(x))$ are uniquely determined such that they are invariant under f , satisfy the matching condition, satisfy equalities (7.5) and (7.6), and are well-defined along the ι -boundaries of the Markov rectangles. Hence, r is an ι -ratio function.*

Proof. The proof follows by induction in $n \geq 0$. Let us prove the case $n = 0$. By construction, $L_i(x) \supset A(f_{\iota}^i(x))$, i.e $L_i(x)$ contains all the ι -boundary leaf segments of Markov rectangles intersecting the global leaf segment passing through the point $f_{\iota}^i(x)$. Hence, if $G_2 \subset L_i(x) \setminus f_{\iota'} L_{i+1}(x)$ is an out-gap of $R(2, i, x)$, then there is an out-gap or a leaf 2-gap segment G_1 of $R(1, i, x)$ such that $G_1 = G_2$. Therefore, we define $r(G_2 : L_i(x)) = r(G_1 : L_i(x))$. Since $L_i(x) \supset A(f_{\iota}^i(x))$, if $G_2 \subset L_i(x) \setminus f_{\iota'} L_{i+1}(x)$ is a leaf 2-gap segment of $R(2, i, x)$ then there is an out-gap or a leaf 2-gap segment G_1 of $R(1, i, x)$ such that $G_1 = G_2$. Hence, we define $r(G_2 : L_i(x)) = r(G_1 : L_i(x))$. If $C_2 \subset L_i(x) \setminus f_{\iota'} L_{i+1}(x)$ is a leaf 2-cylinder segment of $R(2, i, x)$, then there is a primary leaf segment or a leaf 2-cylinder segment C_1 of $R(1, i, x)$ such that $C_2 = C_1$. Therefore, we define $r(C_2 : L_i(x)) = r(C_1 : L_i(x))$. Let us prove that for all out-gaps and $j+2$ -leaf segments $C \subset f_{\iota'}^{n+1} L_i(x) \setminus f_{\iota'}^{n+2} L_{i+1}(x)$ of $f_{\iota'}^{n+1} R(2, i, x)$, with $1 \leq j \leq n+1$, the ratios $r(C : f_{\iota'}^{n+1} L_i(x))$ are uniquely determined using the induction hypotheses with respect to n . By the induction hypotheses and by the matching condition, the ratio $r(f_{\iota} C : f_{\iota'}^n L_i(x))$ is well-defined. By invariance of f , we define $r(C : f_{\iota'}^{n+1} L_i(x)) = r(f_{\iota} C : f_{\iota'}^n L_i(x))$. which ends the proof of the induction.

Lemma 7.7. *Let $(\gamma_1, J_1) \in \mathcal{D}_1(\rho, \delta, P)$. Let $x = x(O)$, where $O \in \mathcal{Q}$, and let $i \in \{0, 1, \dots, p(x)-1\}$. For all out-gaps and all 2-leaf segments $C \subset L_i(x)$ of $R(2, i, x)$, the ratios $r(C : L_i(x))$ are uniquely determined such that they are invariant under f , satisfy the matching condition, satisfy equalities (7.5) and (7.6), and are well-defined along the ι -boundaries of the Markov rectangles. Hence, r is an ι -ratio function.*

Proof. By construction of $L_i(x) \setminus f_{\iota'} L_{i+1}(x)$, there is $k = k(n, i, x)$ such that $L_i(x) \setminus f_{\iota'} L_{i+1}(x) = \cup_{l=1}^k D_l$, where D_l are out-gaps, primary leaf segments and 2-leaf segments of $R(1, i, x)$. Therefore, $f_{\iota'}^n L_i(x) \setminus f_{\iota'}^{n+1} L_{i+1}(x) = \cup_{l=1}^k f_{\iota'}^n D_l$ where $f_{\iota'}^n D_l$ are out-gaps and $j+2$ -leaf segments of $R(1, i, f_{\iota'}^n(x))$ with $0 \leq j \leq n$. Hence, by Lemma 7.6 and using the matching condition, the ratio $r(f_{\iota'}^n L_i(x) \setminus f_{\iota'}^{n+1} L_{i+1}(x) : f_{\iota'}^n L_i(x))$ is well-defined. Hence, using the matching condition, we define

$$r(f_{\iota'}^{n+1} L_{i+1}(x) : f_{\iota'}^n L_i(x)) = 1 - r(f_{\iota'}^n L_i(x) \setminus f_{\iota'}^{n+1} L_{i+1}(x) : f_{\iota'}^n L_i(x)) .$$

Therefore, using again the matching condition, we define

$$(7.11) \quad r(f_{\iota'}^{n+1} L_{i+n+1}(x) : L_i(x)) = \prod_{j=0}^n r(f_{\iota'}^{j+1} L_{i+j+1}(x) : f_{\iota'}^j L_{i+j}(x)) .$$

Let $M(i, x)$ be the 2-cylinder of $R(2, i, x)$ containing the point x . Take $N > 0$, large enough, such that $f_{\iota'}^{N+1} L_{i+N+1}(x) \subset M(i, x)$. Hence, there is $m = m(N, i, x)$ such that $M(i, x) = (\cup_{l=0}^m D_l) \cup f_{\iota'}^{N+1} L_{i+N+1}(x)$ where D_l are out-gaps or $j+2$ -leaf segments of $R(1, i, x)$ for some $0 \leq j \leq N$. Hence, by Lemma 7.6, (7.11) and using the matching condition, the ratio is well-defined by

$$r(M(i, x) : L_i(x)) = \sum_{l=0}^m r(D_l : L_i(x)) + r(f_{\iota'}^{N+1} L_{i+N+1}(x) : L_i(x)) .$$

If $C \subset L_i(x) \setminus M(i, x)$ is a out-gap or a 2-leaf segment of $R(2, i, x)$, then, by Lemma 7.6, the ratio is well-defined by $r(C : L_i(x))$. By construction of the ratios, in Lemmas 7.3-7.7, they are compatible with the cylinder-gap condition.

Definition 7.3. Let $(\gamma_1, J_1) \in \mathcal{D}_1(\rho, \delta, P)$. Let $x = x(O)$, where $O \in \mathcal{Q}$, and let $i \in \{0, 1, \dots, p(x) - 1\}$. Let the ratios $r(C : L_i(x))$ for all out-gaps and all 2-leaf segments $C \subset L_i(x)$ of $R(2, i, x)$ be as given in Lemma 7.7. For all $\xi \in \text{Co}(2, i, x)$, letting $I = f_{i'}\xi \subset L_i(x)$, we define

$$J_2(\xi) = r(I : L_i(x))r(L_i(x) : mI)\rho(I : K_i)^{-1/\delta}e^{-P/\delta}.$$

For all $(C, D) \in \text{Gap}(2, i, x)$, we define

$$\gamma(C : D) = r(f_{i'}C : L_i(x))r(L_i(x) : f_{i'}D).$$

Lemma 7.8. Let $\mathcal{D}_1(\rho, \delta, P) \neq \emptyset$. The gap cocycle map $b = b_{\rho, \delta, P} : \mathcal{D}_1(\rho, \delta, P) \rightarrow \mathcal{D}_2(\rho, \delta, P)$ is well-defined by $b(\gamma_1, J_1) = (\gamma_2, J_2)$ where γ_2 and J_2 are as given in Definition 7.3. Furthermore, the cocycle gap map b is a bijection.

Proof. Let us check that (γ_2, J_2) satisfies properties (i)-(iii) of $\mathcal{D}_2(\rho, \delta, P)$. By construction of the ratios r , in Lemmas 7.3-7.7, (γ_2, J_2) satisfies properties (i) and (ii) in the definition of $\mathcal{D}_2(\rho, \delta, P)$. Let us check property (iii). Let us denote by A and B the $p(x)$ -cylinders of $M(1, x)$ and $M(2, x)$, respectively, containing the point x . By invariance of r , we have that $r(A : B) = r(f_t^{p(x)}A : f_t^{p(x)}B)$, and so $r(A : f_t^{p(x)}A) = r(B : f_t^{p(x)}B)$. By invariance of the ι -measure ratio function ρ , we have that $\rho(A : B) = \rho(f_t^{p(x)}A : f_t^{p(x)}B)$, and so $\rho(A : f_t^{p(x)}A) = \rho(B : f_t^{p(x)}B)$. Since, by hypotheses $\prod_{i=0}^{p(x)-1} J(m^i f_t^{p(x)-i}A) = 1$, we get, from (7.5), that $r(A : f_t^{p(x)}A) = \rho(A : f_t^{p(x)}A)e^{p(x)P/\delta}$. Therefore,

$$\begin{aligned} r(B : f_t^{p(x)}B) &= r(A : f_t^{p(x)}A) \\ &= \rho(A : f_t^{p(x)}A)e^{p(x)P/\delta} \\ &= \rho(B : f_t^{p(x)}B)e^{p(x)P/\delta} \end{aligned}$$

and so, using (7.5), we obtain that $\prod_{i=0}^{p(x)-1} J(\pi_{\mathbf{B}'_0} f_t^i B) = 1$.

Definition 7.4. Let $\mathbf{B}^t$ be a gap train-track. Let $\delta > 0$ and $P \in \mathbb{R}$. Let ρ be an ι -measure ratio function and $\nu = \nu_\rho$ the corresponding Gibbs measure on Θ . We say that a pair (γ, J) is a (ν, δ, P) ι cocycle-gap pair if (γ, J) has the following properties:

- (i) γ is an ι -gap ratio function.
- (ii) J is an ι measure-length ratio cocycle.
- (iii) If $\mathbf{B}^{t'}$ is a no-gap train-track then (γ, J) satisfies the following gap-cocyle property: $b(\gamma|\text{Gap}_1, J|\text{Co}_1) = (\gamma|\text{Gap}_2, J|\text{Co}_2)$, where $b = b_{\nu, \delta, P}$ is the gap-cocyle map.

Let $\mathcal{JG}^t(\nu, \delta, P)$ be the set of all (ν, δ, P) ι cocycle-gap pairs.

Lemma 7.9. Let $\mathbf{B}^t$ be a gap train-track. Let $\delta > 0$ and $P \in \mathbb{R}$. Let ρ be an ι -measure ratio function with corresponding Gibbs measure ν .

- (i) If there is a (ρ, δ, P) ι -measure-length ratio cocycle, then the set $\mathcal{JG}^t(\nu, \delta, P)$ is an infinite dimensional space.
- (ii) If $\mathcal{S}$ is a C^{1+} self-renormalisable structure with natural geometric measure $\mu_{\mathcal{S}, \delta} = i_*\nu$ and pressure P , then $(\gamma_{\mathcal{S}}, J_{\mathcal{S}, \delta}) \in \mathcal{JG}^t(\nu, \delta, P)$.
- (iii) If the set $\mathcal{JG}^t(\nu, \delta, P) \neq \emptyset$, then there is a well-defined injective map $(\gamma, J) \rightarrow r(\gamma, J)$ which associates to each cocycle-gap pair $(\gamma, J) \in \mathcal{JG}^t(\nu, \delta, P)$ an ι -ratio function $r(\gamma, J)$ satisfying (7.5) and (7.6).

Remark 7.10. Let $0 < \delta < 1$ and $P = 0$. Let ρ be an ι -measure ratio function with corresponding Gibbs measure ν . Since $J = 1$ is a (ρ, δ, P) ι -measure-length ratio cocycle, then, by Lemma 7.9, the set $\mathcal{JG}^\iota(\nu, \delta, P)$ is an infinite dimensional space.

Proof of Lemma 7.9. Proof of (i). Choose a map $\gamma_1 : \text{Gap}_1 \rightarrow \mathbb{R}^+$. Let J_0 be a (ρ, δ, P) ι -measure-length ratio cocycle, and let $J_1 = J_0|_{\text{Co}_1}$. Since $(\gamma_1, J_1) \in \mathcal{D}_1(\rho, \delta, P)$, by Lemma 7.8, the pair $(\gamma_2, J_2) = b_{\rho, \delta, P}(\gamma_1, J_1) \in \mathcal{D}_2(\rho, \delta, P)$ is well-defined. Let k_0 and k_2 be maps such that $J_0 = k_0/(k_0 \circ f_\iota)$ and $J_2 = k_2/(k_2 \circ f_\iota)$. For every $x(O) \in \mathcal{Q}$, let $x = x(O)$, and let B be the $p(x)$ -cylinder of $M(2, x)$ containing the point x . Recall that the points $\pi_{\mathbf{B}'_o} B, \pi_{\mathbf{B}'_o} f_\iota B, \dots, \pi_{\mathbf{B}'_o} f_\iota^{p(x)-1} B$ in $\mathbf{B}'_o$ form a periodic orbit under f_ι , with period $p(x)$, and that the primary cylinders contained in the set $\text{Co}(2, i, x)$ are pre-orbits of the points $\pi_{\mathbf{B}'_o} f_\iota^i B$ in $\mathbf{B}'_o$, under f_ι . Therefore, there is a small neighbourhood V of Co_2 in $\mathbf{B}'_o$, there is $\varepsilon > 0$, small enough, and there is an Hölder continuous map $k : \mathbf{B}'_o \rightarrow \mathbb{R}^+$ with the following properties:

- (i) $k|_{\text{Co}_2} = k_2$, $k|_{(\mathbf{B}'_o \setminus V)} = k_0$ and $\text{Co}_1 \subset \mathbf{B}'_o \setminus V$.
- (ii) Let $a = \min_{\xi \in \text{Co}_2} \{J_0(\xi), J_2(\xi)\}$ and $b = \max_{\xi \in \text{Co}_2} \{J_0(\xi), J_2(\xi)\}$, and let $J = k/(k \circ f_\iota)$. For every $\xi \in V$, we have that $a - \varepsilon \leq J(\xi) \leq b + \varepsilon$, and, so, J satisfies the cocycle-gap property.

Choosing an Hölder continuous map $\gamma : \mathcal{G}' \rightarrow \mathbb{R}^+$ such that $\gamma|_{\text{Gap}_1} = \gamma_1$ and $\gamma|_{\text{Gap}_2} = \gamma_2$ and by property (i) above, the pair (γ, J) satisfies (7.1). Therefore, the pair (γ, J) is contained in $\mathcal{JG}^\iota(\nu, \delta, P)$. Using that (7.1) is an open condition, the above construction allows us to construct an infinite set of ι -measure-length ratio cocycles and an infinite set of gap ratio functions such that the corresponding pairs are contained in $\mathcal{JG}^\iota(\nu, \delta, P)$.

Proof of (ii). Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure with natural geometric measure $\mu_{\mathcal{S}, \delta} = i_* \nu$ and pressure P . By Lemma 7.1, $J_{\mathcal{S}, \delta}$ is a (ρ, δ, P) ι -measure-length ratio cocycle and, by (7.4), $\gamma_{\mathcal{S}}$ is an ι -gap ratio function. If $\mathbf{B}'$ is a no-gap train-track, using (7.5) and (7.6), the pair $(\gamma_{\mathcal{S}}, J_{\mathcal{S}, \delta})$ satisfies the cocycle-gap condition because the ratio function $r_{\mathcal{S}}$ associated to $\mathcal{S}$ is well-defined along the ι -boundaries of the Markov rectangles.

Proof of (iii). The equations (7.5) and (7.6) give us an inductive construction, on the level n of the n -cylinders and n -gaps, of a ratio function r in terms of $(\rho, J, \gamma, \delta, P)$ with the property that the ratio between a leaf n -cylinder segment C and a leaf n -cylinder or n -gap segment D with a common endpoint with C is bounded away from zero and infinity independent of n and of the cylinders and gaps considered. The construction gives that r is invariant under f . The Hölder continuity of γ , J and ρ implies that r satisfies (3.2). If $\mathbf{B}'$ is a no-gap train-track, by the construction of the cocycle-gap condition, the ratio function r is well-defined along the ι -boundaries of the Markov rectangles. Hence, r is an ι -ratio function. $\square$

8. REALISATIONS OF GIBBS MEASURES

In this section, we are going to give an explicit geometric construction of all C^{1+} hyperbolic diffeomorphisms which have a natural geometric measure, and we will prove the theorems stated in the introduction of the paper.

8.1. One-dimensional realisations. Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure on a train-track $\mathbf{B}^\iota$. In Lemma 6.8 we have shown that the map

$$(8.1) \quad (\mathcal{S}, \delta) \rightarrow \rho_{\mathcal{S}, \delta}$$

is well-defined where $\rho_{\mathcal{S}, \delta}$ is the ι -measure ratio function associated to a Gibbs measure $\nu_{\mathcal{S}, \delta} = \nu$ such that $\mu_{\mathcal{S}, \delta} = (i_\iota)_* \nu_\iota$ is a natural geometric measure of $\mathcal{S}$.

Lemma 8.1. (*Rigidity*) Let $\mathbf{B}^\iota$ be a no gap train-track (and $\delta = 1$). The map $\mathcal{S} \rightarrow \rho_{\mathcal{S},\delta}$ is an one-to-one correspondence between C^{1+} self-renormalisable structures on $\mathbf{B}^\iota$ and ι -measure ratio functions. Furthermore, $\rho_{\mathcal{S},\delta} = r_{\mathcal{S}}$ where $r_{\mathcal{S}}$ is the ratio function determined by $\mathcal{S}$.

However, if $\mathbf{B}^\iota$ is a gap train-track then the set of pre-images of the map $(\mathcal{S}, \delta) \rightarrow \rho_{\mathcal{S},\delta}$ is an infinite dimensional space (see Lemma 8.3 below).

Proof. By Theorem 6.1, the C^{1+} self-renormalisable structure $\mathcal{S}$ realises a Gibbs measure $\nu = \nu_{\mathcal{S},\delta}$. By Lemma 6.8, we get that $\rho_{\mathcal{S},\delta} = r_{\mathcal{S}}$. Since, by Lemma 4.1, the ratio function $r_{\mathcal{S}}$ determines uniquely the C^{1+} self-renormalisable structure $\mathcal{S}$, the map $\mathcal{S} \rightarrow \rho_{\mathcal{S},\delta}$ is an one-to-one correspondence.

Definition 8.1. Let $\mathbf{B}_\iota$ and $\mathbf{B}_{\iota'}$ be (gap or no-gap) train-tracks. Let ρ be an ι -measure ratio function and $\nu = \nu_\rho$ on Θ the corresponding Gibbs measure (see Remark 5.8). Let us denote by $\mathcal{D}^\iota(\nu, \delta, P)$ the set of all C^{1+} self-renormalisable structures $\mathcal{S}$ with geometric natural measure $\mu_{\mathcal{S},\delta} = (i_\iota)_* \nu_\iota$ and pressure P .

By Lemma 8.1, if $\mathbf{B}^\iota$ is a no-gap train-track, and $\delta = 1$ and $P = 0$, the set $\mathcal{D}^\iota(\nu, \delta, P)$ is a singleton.

Let $\mathbf{B}^\iota$ be a gap train-track and $\mathcal{S}$ a C^{1+} self-renormalisable structure in $\mathcal{D}^\iota(\nu, \delta, P)$. In Theorem 7.2, we associate to the C^{1+} self-renormalisable structure $\mathcal{S}$ a measure-length ratio cocycle $J_{\mathcal{S}}$, and, in Section 7.1, we associate to the C^{1+} self-renormalisable structure $\mathcal{S}$ a gap ratio function $\gamma_{\mathcal{S}}$. By Lemma 7.9, if $\mathbf{B}^{\iota'}$ is a no-gap train-track then the cylinder-gap condition of $r_{\mathcal{S}}$ implies that the pair $(\gamma_{\mathcal{S}}, J_{\mathcal{S},\delta})$ satisfies the cocycle-gap condition. Therefore, the map

$$(8.2) \quad \mathcal{S} \rightarrow (\gamma_{\mathcal{S}}, J_{\mathcal{S},\delta})$$

between C^{1+} self-renormalisable structures contained in $\mathcal{D}^\iota(\nu, \delta, P)$ and pairs contained in $\mathcal{JG}^\iota(\nu, \delta, P)$ is well-defined.

Definition 8.2. The (δ_ι, P_ι) -bounded solenoid equivalence class of a Gibbs measure ν is the set of all solenoid functions σ_ι with the following properties: There is $C = C(\sigma_\iota) > 0$ such that for every pair $(\xi, D) \in \text{msc}_\iota$

$$|\delta_\iota \log s_\iota(D_\Lambda \cap \xi_\Lambda : \xi_\Lambda) - \log \rho_\xi(D) - nP_\iota| < C,$$

where (i) ρ is the ι -extended measure scaling function of ν , (ii) s_ι is the scaling function determined by σ_ι , (iii) $\xi_\Lambda = i(\pi_{\iota'}^{-1}\xi)$ is an ι' -leaf primary cylinder segment and (iv) $D_\Lambda = i(\pi_\iota^{-1}D)$ and so $D_\Lambda \cap \xi_\Lambda$ is an ι -leaf n -cylinder segment.

Remark 8.2. Let $\sigma_{1,\iota}$ and $\sigma_{2,\iota}$ be two solenoid functions in the same (δ_ι, P_ι) -bounded solenoid equivalence class of a Gibbs measure ν . Using the fact that $\sigma_{1,\iota}$ and $\sigma_{2,\iota}$ are bounded away from zero, we obtain that the corresponding scaling functions also satisfy inequality (3.6) for all pairs $(J, m^i J)$ where J is an ι -leaf $(i+1)$ -cylinder. Hence, the solenoid functions $\sigma_{1,\iota}$ and $\sigma_{2,\iota}$ are in the same bounded equivalence class (see Definiton 3.2).

By Lemma 8.3, below, the set $\mathcal{JG}^\iota(\nu, \delta, P)$ gives a parametrization of all C^{1+} self-renormalisable structures $\mathcal{S}$ which are pre-images of the ι -measure ratio function $\rho_{\nu,\iota}$, under the map $\mathcal{S} \rightarrow \rho_{\mathcal{S},\delta}$ given in (8.1), with a natural geometric measure $\mu_\iota = (i_\iota)_* \nu_\iota$ and pressure $P(\mathcal{S}, \delta) = P$. Hence, $\mathcal{JG}^\iota(\nu, \delta, P)$ forms a moduli space for the set of all C^{1+} self-renormalisable structures in $\mathcal{D}^\iota(\nu, \delta, P)$.

Lemma 8.3. (*Flexibility*) Let $\mathbf{B}_\iota$ be a gap train-track. Let ρ be an ι -measure ratio function and $\nu = \nu_\rho$ the corresponding Gibbs measure on Θ .

- (i) Let $\delta > 0$ and $P \in \mathbb{R}$ be such that $\mathcal{JG}^\iota(\nu, \delta, P) \neq \emptyset$. The map $\mathcal{S} \rightarrow (\gamma_{\mathcal{S}}, J_{\mathcal{S},\delta})$ determines a one-to-one correspondence between C^{1+} self-renormalisable structures in $\mathcal{D}^\iota(\nu, \delta, P)$ and cocycle-gap pairs in $\mathcal{JG}^\iota(\nu, \delta, P)$.

(ii) A C^{1+} self-renormalisable structure $\mathcal{S}$ is contained in $\mathcal{D}^\iota(\nu, \delta, P)$ if, and only if, the ι -solenoid function $\sigma_{\mathcal{S}}$ is contained in the (δ, P) -bounded solenoid equivalence class of ν (see Definition 8.2).

Proof of (i). Let us prove that $(J, \gamma) \in \mathcal{JG}^\iota(\nu, \delta, P)$ determines a unique C^{1+} self-renormalisable structure $\mathcal{S}$ with a natural geometric measure $\mu_{\mathcal{S}, \delta} = (i_\iota)_* \nu_\iota$. By Lemma 7.9, the pair (J, γ) determines a unique ι -ratio function $r = r_\iota(J, \gamma)$. By Lemma 4.1, the ι -ratio function r determines a unique C^{1+} self-renormalisable structure $\mathcal{S}$ with an atlas $\mathcal{B}(r)$. Let us prove that $\mu_\iota = (i_\iota)_* \nu_\iota$ is a natural geometric measure of $\mathcal{S}$ with the given δ and P . Let ρ be the ι -measure ratio function associated to the Gibbs measure ν . By Lemma 5.2, for every leaf n -cylinder or n -gap segment I we obtain that

$$(8.3) \quad \mu_\iota(I) = \mathcal{O}(\rho(I \cap \xi : \xi))$$

for every $\xi \in \pi_{\iota'}(I)$. On the other hand, by construction of the ratio function r_ι and using (7.5), we get

$$\rho(I \cap \xi : \xi) = e^{-nP} r(I \cap \xi : \xi)^\delta \prod_{j=0}^{n-1} \left(J \left(\tau_{\iota'}^j(\xi) \right) \right)^{-\delta}.$$

Since J is a Hölder cocycle, it follows that $\prod_{j=0}^{n-1} J \left(\tau_{\iota'}^j(\xi) \right) = k(\xi)/k(\tau_{\iota'}^n(\xi))$ is uniformly bounded away from zero and infinity, where k is an Hölder continuous positive function. By (4.1), we get that

$$(8.4) \quad r(I \cap \xi : \xi) = \mathcal{O}(|I|_j)$$

where $j \in \mathcal{B}(r)$ and I is contained in the domain of j . Hence,

$$(8.5) \quad \rho(I \cap \xi : \xi) = \mathcal{O}(|I|_j^\delta e^{-nP}).$$

Putting together equations (8.3) and (8.5), we deduce that $\mu_\iota(I) = \mathcal{O}(|I|_j^\delta e^{-nP})$, and so $\mu_\iota = (i_\iota)_* \nu_\iota$ is a natural geometric measure of $\mathcal{S}$ with the given δ and P .

Proof of (ii). Let $\mathcal{S}$ be a C^{1+} self-renormalisable structure in $\mathcal{D}^\iota(\nu, \delta, P)$. Then, putting together (8.4) and (8.5), there is $\kappa > 0$ such that

$$|\delta \log r_\iota(I \cap \xi : \xi) - \log \rho(I \cap \xi : \xi) - np| < \kappa$$

for every leaf n -cylinder I and $\xi \in \pi_{\iota'}(I)$. Thus the solenoid function $r|\mathbf{S}^\iota$ is in the (δ, P) -bounded solenoid equivalence class of ν .

Conversely, let $\mathcal{S}$ be a C^{1+} self-renormalisable structure in the (δ, P) -bounded solenoid equivalence class of ν and $\mu_\iota = (i_\iota)_* \nu_\iota$, i.e. there is $\kappa > 0$ such that

$$(8.6) \quad |\delta \log r_\iota(I \cap \xi : \xi) - \log \rho(I \cap \xi : \xi) - np| < \kappa$$

for every leaf n -cylinder I and $\xi \in \pi_{\iota'}(I)$. Hence, using (8.3) and (8.4) in (8.6), we get $\mu_\iota(I) = \mathcal{O}(|I|_j^\delta e^{-nP})$. Since $\mu_\iota = (i_\iota)_* \nu_\iota$ we get that $\mathcal{S}$ is contained in $\mathcal{D}^\iota(\nu, \delta, P)$. $\square$

8.2. Two-dimensional realisations. We start by giving the definition of a natural geometric measure for a C^{1+} hyperbolic diffeomorphism.

Definition 8.3. For $\iota \in \{s, u\}$, if $\mathbf{B}^\iota$ is a gap train-track assume $0 < \delta_\iota < 1$, and if $\mathbf{B}^\iota$ is a no-gap train-track take $\delta_\iota = 1$.

(i) Let g be a C^{1+} hyperbolic diffeomorphism in $\mathcal{T}(f, \Lambda)$. We say that g has a natural geometric measure $\mu = \mu_{g, \delta_s, \delta_u}$ with pressures $P_s = P_s(g, \delta_s, \delta_u)$ and $P_u = P_u(g, \delta_s, \delta_u)$ if, there is $\kappa > 1$ such that for all leaf n_s -cylinder I_s , for all leaf n_u -cylinder I_u ,

$$(8.7) \quad \kappa^{-1} < \frac{\mu(R)}{|I_u|^{\delta_u} |I_s|^{\delta_s} e^{-n_s P_s - n_u P_u}} < \kappa,$$

where R is the (n_s, n_u) -rectangle $[I_s, I_u]$ and where the lengths $|\cdot|$ are measured in the stable and unstable C^{1+} foliated lamination atlases $\mathcal{A}_s(g, \rho)$ and $\mathcal{A}_u(g, \rho)$ of g with respect to some Riemannian metric ρ .

(ii) We say that a C^{1+} hyperbolic diffeomorphism with a natural geometric measure $\mu = \mu_{g, \delta_s, \delta_u}$ with pressures $P_s = P_s(g, \delta_s, \delta_u)$ and $P_u = P_u(g, \delta_s, \delta_u)$ is a C^{1+} realisation of a Gibbs measure $\nu = \nu_{g, \delta_s, \delta_u}$ if $\mu = i_* \nu$. We denote by $\mathcal{T}(\nu, \delta_s, P_s, \delta_u, P_u)$ the set of all these C^{1+} hyperbolic diffeomorphisms $g \in \mathcal{T}(f, \Lambda)$.

A C^{1+} hyperbolic diffeomorphism $g \in \mathcal{T}(f, \Lambda)$ determines an unique pair $(\mathcal{S}(g, s), \mathcal{S}(g, u))$ of C^{1+} stable and unstable self-renormalisable structures (see Lemma 4.2). By Theorem 6.1, for $\delta_s > 0$ and $\delta_u > 0$, the pair $(\mathcal{S}(g, s), \mathcal{S}(g, u))$ of self-renormalisable structures determines an unique pair of natural geometric measures $(\mu_{\mathcal{S}(g, s), \delta_s}, \mu_{\mathcal{S}(g, u), \delta_u})$ corresponding to a unique pair of Gibbs measures $(\nu_{\mathcal{S}(g, s), \delta_s}, \nu_{\mathcal{S}(g, u), \delta_u})$. Furthermore, by Lemma 6.8, the self-renormalisable structures $(\mathcal{S}(g, s), \mathcal{S}(g, u))$ determine a pair of measure ratio functions $(\rho_{\mathcal{S}(g, s), \delta_s}, \rho_{\mathcal{S}(g, u), \delta_u})$ of $(\nu_{\mathcal{S}(g, s), \delta_s}, \nu_{\mathcal{S}(g, u), \delta_u})$.

Lemma 8.4. *Let g be a C^{1+} hyperbolic diffeomorphism contained in $\mathcal{T}(f, \Lambda)$. The following statements are equivalent:*

- (i) g has a natural geometric measure $\mu_{g, \delta_s, \delta_u}$;
- (ii) g is a C^{1+} realisation of a Gibbs measure $\nu_{g, \delta_s, \delta_u}$;
- (iii) $\nu_{\mathcal{S}(g, s), \delta_s} = \nu_{\mathcal{S}(g, u), \delta_u}$;
- (iv) The s -measure ratio function $\rho_{\mathcal{S}(g, s), \delta_s}$ is dual to the u -measure ratio function $\rho_{\mathcal{S}(g, u), \delta_u}$.

Furthermore, if g has a natural geometric measure $\mu_{g, \delta_s, \delta_u}$, then $(\pi_s)_* \mu_{g, \delta_s, \delta_u} = \mu_{\mathcal{S}(g, s), \delta_s}$ and $(\pi_u)_* \mu_{g, \delta_s, \delta_u} = \mu_{\mathcal{S}(g, u), \delta_u}$.

Proof. By Lemma 6.9, (iii) is equivalent to (iv). By definition if g is a C^{1+} realisation of a Gibbs measure $\nu_{g, \delta_s, \delta_u}$ then $\mu_{g, \delta_s, \delta_u} = i_* \nu_{g, \delta_s, \delta_u}$ is a natural geometric measure of g , and so (ii) implies (i). Let us prove first that (i) implies (ii) and (iii), and secondly that (iii) implies (i). Then the last paragraph of this lemma follows from (8.10) below which ends the proof.

(i) implies (ii) and (iii). Let $\mu_{g, \delta_s, \delta_u}$ be the natural geometric measure of g . Since the stable and unstable lamination atlases $\mathcal{A}_s(g, \rho)$ and $\mathcal{A}_u(g, \rho)$ of g are C^{1+} foliated (see Section 2.13) and by construction of the C^{1+} train-track atlases $\mathcal{B}_s(g, \rho)$ and $\mathcal{B}_u(g, \rho)$, in Section 4.2, we obtain that there is $\kappa_1 \geq 1$ with the property that, (for $\iota = s$ and u) and for every ι -leaf n_ι -cylinder I ,

$$(8.8) \quad \kappa_1^{-1} |I|_\rho \leq |I'|_j \leq \kappa_1 |I|_\rho$$

where $I' = \pi_{\mathbf{B}^\iota}(I)$, where $|I'|_j$ is measured in any chart $j \in \mathcal{B}_\iota(g, \rho)$ and where $|I|_\rho$ is the length in the Riemannian metric ρ of the minimal full ι -leaf containing I . Let I'_Λ be the $(1, n_\iota)$ -rectangle in Λ such that $\pi_{\mathbf{B}^\iota}(I'_\Lambda) = I'$. Noting that $(\pi_{\mathbf{B}^\iota})_* \mu_{g, \delta_s, \delta_u}(I') = \mu_{g, \delta_s, \delta_u}(I'_\Lambda)$, by (8.7) and (8.8), there is $\kappa_2 \geq 1$ such that

$$(8.9) \quad \kappa_2^{-1} \leq \frac{(\pi_{\mathbf{B}^\iota})_* \mu_{g, \delta_s, \delta_u}(I')}{|I'|_j^{\delta_\iota} e^{-n_\iota P_\iota}} \leq \kappa_2,$$

for every n_ι -cylinder I' on the train-track. By Theorem 6.1, the natural geometric measure determined by the C^{1+} self-renormalisable structure $\mathcal{S}(g, \iota)$ and by $\delta_\iota > 0$ is uniquely determined by (8.9). Hence,

$$(8.10) \quad (\pi_{\mathbf{B}^s})_* \mu_{g, \delta_s, \delta_u} = \mu_{\mathcal{S}(g, s), \delta_s} \text{ and } (\pi_{\mathbf{B}^u})_* \mu_{g, \delta_s, \delta_u} = \mu_{\mathcal{S}(g, u), \delta_u} .$$

Therefore, the Gibbs measures $\nu_{\mathcal{S}(g, s), \delta_s}$ and $\nu_{\mathcal{S}(g, u), \delta_u}$ on Θ are equal which proves (iii), and $\mu_{g, \delta_s, \delta_u} = i_* \nu_{\mathcal{S}(g, s), \delta_s} = i_* \nu_{\mathcal{S}(g, u), \delta_u}$ which proves (ii).

(iii) implies (i). Let us denote $\nu_{\mathcal{S}(g, s), \delta_s} = \nu_{\mathcal{S}(g, u), \delta_u}$ by ν . Let $\mu = i_* \nu$. For $\iota \in \{s, u\}$, by definition of a C^{1+} realisation of a Gibbs measure as a self-renormalisable structure $\mathcal{S}(g, \delta_\iota)$, for every ι -leaf n_ι -cylinder I_ι , there is $\kappa_3 \geq 1$ such that

$$\kappa_3^{-1} \leq \frac{\mu_\iota(I'_\iota)}{|I'_\iota|_{\delta_\iota} e^{-n_\iota P_\iota}} \leq \kappa_3 ,$$

where $I'_\iota = \pi_{\mathbf{B}^\iota}(I)$ and $|I'_\iota|_j$ is measured in any chart $j \in \mathcal{B}_\iota(g, \rho)$. Hence, by (8.8), for $\iota = s$ and u , we obtain that

$$(8.11) \quad \mu_\iota(I'_\iota) = \mathcal{O}\left(|I_\iota|_\rho^{\delta_\iota} e^{-n_\iota P_\iota}\right) .$$

Let R be the rectangle $[I_s, I_u]$. By Theorem 6.4,

$$\mu(R) = \int_{I'_u} \rho_{\iota, \xi}(R : M) \mu_{\iota'}(d\xi) ,$$

where M is the Markov rectangle containing R . By Theorem 6.1 (i) and (ii), we get that $\rho_{\iota, \xi}(R : M) = \mathcal{O}(\mu_\iota(I'_\iota))$ for every $\xi \in \pi_{\mathbf{B}^{\iota'}}(R)$. Hence

$$(8.12) \quad \mu(R) = \mathcal{O}(\mu_s(I'_s) \mu_u(I'_u)) .$$

Putting together (8.11) and (8.12), we get

$$(8.13) \quad \mu(R) = \mathcal{O}\left(|I_u|_\rho^{\delta_u} |I_s|_\rho^{\delta_s} e^{-n_s P_s - n_u P_u}\right)$$

and so μ is a natural geometric measure.

Lemma 8.5. *The map $g \rightarrow (\mathcal{S}(g, s), \mathcal{S}(g, u))$ gives a one-to-one correspondence between C^{1+} conjugacy classes of hyperbolic diffeomorphisms contained in $\mathcal{T}(\nu, \delta_s, P_s, \delta_u, P_u)$ and pairs of C^{1+} self-renormalisable structures contained in $\mathcal{D}^s(\nu, \delta_s, P_s) \times \mathcal{D}^u(\nu, \delta_u, P_u)$.*

Proof. By Lemma 8.4, if $g \in \mathcal{T}(\nu, \delta_s, P_s, \delta_u, P_u)$ then, for $\iota \in \{s, u\}$, $\mathcal{S}(g, \iota) \in \mathcal{D}^\iota(\nu, \delta_\iota, P_\iota)$. Conversely, by Lemma 4.2, a pair $(\mathcal{S}_s, \mathcal{S}_u) \in \mathcal{D}^s(\nu, \delta_s, P_s) \times \mathcal{D}^u(\nu, \delta_u, P_u)$ determines a C^{1+} hyperbolic diffeomorphism g such that $\mathcal{S}(g, s) = \mathcal{S}_s$ and $\mathcal{S}(g, u) = \mathcal{S}_u$ and $\nu_{\mathcal{S}(g, s), \delta_s} = \nu_{\mathcal{S}(g, u), \delta_u} = \nu$. Therefore, by Lemma 8.4, we obtain that g is a C^{1+} realisation of the Gibbs measure ν .

Lemma 8.6. *(Dual-rigidity) Let $\mathbf{B}^{\iota'}$ be a no-gap train-track (and so $\delta_{\iota'} = 1$ and $P_{\iota'} = 0$). For every $\delta_\iota > 0$ and every C^{1+} ι -self-renormalisable structure $\mathcal{S}_\iota$ there is an unique C^{1+} ι' -self-renormalisable structure $\mathcal{S}_{\iota'}$ such that the C^{1+} hyperbolic diffeomorphism g corresponding to the pair $(\mathcal{S}_s, \mathcal{S}_u) = (\mathcal{S}(g, s), \mathcal{S}(g, u))$ has a natural geometric measure $\mu_{g, \delta_s, \delta_u}$. Furthermore, $\mu_{\mathcal{S}_s, \delta_s} = (\pi_{\mathbf{B}^s})_* \mu_{g, \delta_s, \delta_u}$ and $\mu_{\mathcal{S}_u, \delta_u} = (\pi_{\mathbf{B}^u})_* \mu_{g, \delta_s, \delta_u}$.*

Proof. By Theorem 6.1, a C^{1+} self-renormalisable structure $\mathcal{S}_\iota$ and $\delta_\iota > 0$ determine an unique Gibbs measure $\nu = \nu_{\mathcal{S}_\iota, \delta_\iota}$ and $P_\iota \in \mathbb{R}$ such that $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota, P_\iota)$ is a C^{1+} realisation of ν . By Lemma 6.8, the C^{1+} self-renormalisable structure $\mathcal{S}_\iota$ determines an ι -measure ratio function $\rho_{\mathcal{S}_\iota, \delta_\iota}$ for the Gibbs measure ν . By Lemma 6.9, the ι -measure ratio function $\rho_{\mathcal{S}_\iota, \delta_\iota}$ determines an unique ι' -measure ratio function $\rho_{\iota'}$ of ν on Θ . By Lemma 8.1, there is an unique C^{1+} self-renormalisable

structure $\mathcal{S}_{\iota'}$, with ι' -measure ratio function $\rho_{\mathcal{S}_{\iota'},1} = \rho_{\iota'}$, which is a C^{1+} realisation of the Gibbs measure ν . By Lemma 4.2, the pair $(\mathcal{S}_s, \mathcal{S}_u)$ determines a C^{1+} hyperbolic diffeomorphism g such that $\mathcal{S}(g, s) = \mathcal{S}_s$ and $\mathcal{S}(g, u) = \mathcal{S}_u$. Hence, $\nu_{\mathcal{S}(g,s),\delta_s} = \nu$ and $\nu_{\mathcal{S}(g,u),\delta_u} = \nu$ which implies that $\nu_{\mathcal{S}(g,s),\delta_s} = \nu_{\mathcal{S}(g,u),\delta_u}$. Therefore, by Lemma 8.4, g is a C^{1+} realisation of the Gibbs measure ν with natural geometric measure $\mu_{g,\delta_s,\delta_u} = i_*\nu$. Thus, $\mu_{\mathcal{S}_s,\delta_s} = (\pi_{\mathbf{B}^s})_*\mu_{g,\delta_s,\delta_u}$ and $\mu_{\mathcal{S}_u,\delta_u} = (\pi_{\mathbf{B}^u})_*\mu_{g,\delta_s,\delta_u}$.

Recall the definition of the maps $g \rightarrow (\mathcal{S}(g, s), \mathcal{S}(g, u))$ and $\mathcal{S}(g, \iota) \rightarrow (\gamma_{\mathcal{S}(g,\iota)}, J_{\mathcal{S}(g,\iota),\delta_\iota})$ for ι equal to s and u .

Theorem 8.7. (*Flexibility*) Let $\mathbf{B}_\iota$ be a gap train-track. Let ν be a Gibbs measure determining an ι -measure ratio function. Let $\delta_\iota > 0$ and $P_\iota \in \mathbb{R}$ be such that $\mathcal{JG}^\iota(\nu, \delta_\iota, P_\iota) \neq \emptyset$.

(i) (*Smale horseshoes*) Let $\delta_{\iota'} > 0$ and $P_{\iota'} \in \mathbb{R}$ be such that $\mathcal{JG}^{\iota'}(\nu, \delta_{\iota'}, P_{\iota'}) \neq \emptyset$. The map

$$g \rightarrow (\gamma_{\mathcal{S}(g,s)}, J_{\mathcal{S}(g,s),\delta_s}, \gamma_{\mathcal{S}(g,u)}, J_{\mathcal{S}(g,u),\delta_u})$$

gives an one-to-one correspondence between C^{1+} conjugacy classes of hyperbolic diffeomorphisms in $\mathcal{T}(\nu, \delta_s, P_s, \delta_u, P_u)$ and pairs of stable and unstable cocycle-gap pairs in $\mathcal{JG}^s(\nu, \delta_s, P_s) \times \mathcal{JG}^u(\nu, \delta_u, P_u)$.

(ii) (*Codimension one attractors and repellers*) Let $\delta_{\iota'} = 1$ and $P_{\iota'} = 0$. The map $g \rightarrow (\gamma_{\mathcal{S}(g,\iota)}, J_{\mathcal{S}(g,\iota),\delta_\iota})$ gives an one-to-one correspondence between C^{1+} conjugacy classes of hyperbolic diffeomorphisms in $\mathcal{T}(\nu, \delta_s, P_s, \delta_u, P_u)$ and pairs of stable and unstable cocycle-gap pairs in $\mathcal{JG}^\iota(\nu, \delta_\iota, P_\iota)$.

Proof. Statement (i) follows from putting together the results of lemmas 8.3 and 8.5. Statement (ii) follows as statement (i) using the fact that, by Lemma 8.1, the C^{1+} self-renormalisable structure $\mathcal{S}(g, \iota)$ uniquely determines $\mathcal{S}(g, \iota')$ in this case.

9. EIGENVALUES

In this section, we show that the set of stable and unstable eigenvalues of all periodic points of the hyperbolic diffeomorphisms is a complete invariant of the Lipschitz conjugacy classes extending the results of De la Llave, Marco and Moriyo. Furthermore, we extend the eigenvalue formula of A. N. Livšic and Ja. G. Sinai from Anosov diffeomorphisms to C^{1+} hyperbolic diffeomorphisms.

9.1. Lipschitz conjugacy classes.

Lemma 9.1. Let $\mathbf{B}^\iota$ be a (gap or a no-gap) train-track. Let $\delta > 0$ and $P \in \mathbb{R}$. Let $\mathcal{S}_1 \in \mathcal{D}^\iota(\nu_1, \delta, P)$ and $\mathcal{S}_2 \in \mathcal{D}^\iota(\nu_2, \delta, P)$ be C^{1+} self-renormalisable structures. The following statements are equivalent:

- (i) The C^{1+} self-renormalisable structures $\mathcal{S}_1$ and $\mathcal{S}_2$ are Lipschitz conjugate;
- (ii) The Gibbs measures ν_1 and ν_2 are equal;
- (iii) The solenoid functions $s_{\mathcal{S}_1}$ and $s_{\mathcal{S}_2}$ are in the same bounded equivalence class (Definition 3.2).

Proof that (i) is equivalent to (ii). Using (6.1), if $\nu_1 = \nu_2$ then the C^{1+} self-renormalisable structure $\mathcal{S}_1$ is Lipschitz conjugate to $\mathcal{S}_2$. Conversely, if $\mathcal{S}_1$ is Lipschitz conjugate to $\mathcal{S}_2$ then the C^{1+} self-renormalisable structure $\mathcal{S}_1$ (and $\mathcal{S}_2$) satisfies (6.1) with respect to the measures $\mu_{\iota,1} = (i_\iota)_*\nu_{\iota,1}$ and $\mu_{\iota,2} = (i_\iota)_*\nu_{\iota,2}$. By Theorem 6.1, there is a unique τ -invariant Gibbs measure satisfying (6.1) and so $\nu_1 = \nu_2$.

Proof that (ii) is equivalent to (iii). Using (3.6) and (4.1), we obtain that the C^{1+} self-renormalisable structures $\mathcal{S}$ and $\mathcal{S}'$ on $\mathbf{B}^\iota$ are in the same Lipschitz equivalence class if, and only if, the corresponding solenoid functions $r_{\mathcal{S},\iota}|\mathbf{S}^\iota$ and $r_{\mathcal{S}',\iota}|\mathbf{S}^\iota$ are in the same bounded equivalence class. Hence, statement (ii) is equivalent to statement (iii). $\square$




FIGURE 16. An orthogonal chart.

Lemma 9.2. *Let g_1 and g_2 be C^{1+} hyperbolic diffeomorphisms. The following statements are equivalent:*

- (i) *The diffeomorphism g_1 is Lipschitz conjugate to g_2 .*
- (ii) *For ι equal to s and u , $\mathcal{S}(g_1, \iota)$ is Lipschitz conjugate to $\mathcal{S}(g_2, \iota)$.*
- (iii) *For ι equal to s and u , the solenoid functions $s_{g_1, \iota}$ and $s_{g_2, \iota}$ are in the same bounded equivalence class (Definition 3.2).*

Proof that (i) is equivalent to (ii). For all $x \in \Lambda$, let A be a small open set of M containing x , and let R be a rectangle (not necessarily a Markov rectangle) such that $A \cap \Lambda \subset R$. We construct an orthogonal chart $j : R \rightarrow \mathbb{R}^2$ as follows. Let $e_{g,s} : \ell^s(x, R) \rightarrow \mathbb{R}$ be a chart contained in $\mathcal{A}^s(g, \rho)$ and $e_{g,u} : \ell^u(x, R) \rightarrow \mathbb{R}$ be a chart contained in $\mathcal{A}^u(g, \rho)$. The orthogonal chart j on R is now given by $j(z) = (e_{g,s}[z, x], e_{g,u}[x, z]) \in \mathbb{R}^2$ (see Figure 16). By [24], the orthogonal chart $j : R \rightarrow \mathbb{R}^2$ has an extension $\hat{j} : B \rightarrow \mathbb{R}^2$ to an open set B of the surface such that $\hat{j}$ is C^{1+} compatible with the charts in the C^{1+} structure $\mathcal{C}(g)$ of the surface M . Hence, using the orthogonal charts, any two C^{1+} hyperbolic diffeomorphisms g_1 and g_2 are Lipschitz conjugate if, and only if the charts in $\mathcal{A}^\iota(g_1, \rho_1)$ are bi-Lipschitz compatible with the charts in $\mathcal{A}^\iota(g_2, \rho_2)$ for ι equal to s and u . By construction of the train-track atlases $\mathcal{B}^\iota(g_1, \rho_1)$ and $\mathcal{B}^\iota(g_2, \rho_2)$ from the lamination atlases $\mathcal{A}^\iota(g_1, \rho_1)$ and $\mathcal{A}^\iota(g_2, \rho_2)$, the charts in $\mathcal{A}^\iota(g_1, \rho_1)$ are bi-Lipschitz compatible with the charts in $\mathcal{A}^\iota(g_2, \rho_2)$ if, and only if, the charts in $\mathcal{B}^\iota(g_1, \rho_1)$ are bi-Lipschitz compatible with the charts in $\mathcal{B}^\iota(g_2, \rho_2)$. Hence, the C^{1+} hyperbolic diffeomorphisms g_1 and g_2 are Lipschitz conjugate if, and only if, for ι equal to s and u , the corresponding C^{1+} self-renormalisable structures $\mathcal{S}(g_1, \iota)$ and $\mathcal{S}(g_2, \iota)$ are Lipschitz conjugate. Therefore, statement (i) is equivalent to statement (ii).

Proof that (ii) is equivalent to (iii). Follows from Lemma 9.1. $\square$

Lemma 9.3. *Let $\delta_s > 0$, $\delta_u > 0$ and $P_s, P_u \in \mathbb{R}$.*

- (i) *A C^{1+} hyperbolic diffeomorphism g is contained in $\mathcal{T}(\nu, \delta_s, P_s, \delta_u, P_u)$ if, and only if, for ι equal to s and u , the ι -solenoid function $\sigma_{g, \iota}$ is contained in the (δ_ι, P_ι) -bounded solenoid equivalence class of ν (see Definition 8.2).*
- (ii) *If $g_1 \in \mathcal{T}(\nu_1, \delta_s, P_s, \delta_u, P_u)$ and $g_2 \in \mathcal{T}(\nu_2, \delta_s, P_s, \delta_u, P_u)$ are C^{1+} hyperbolic diffeomorphisms then g_1 is Lipschitz conjugate to g_2 if, and only if, $\nu_1 = \nu_2$.*

Proof of (i). By Lemma 4.2, the C^{1+} hyperbolic diffeomorphism g determines a unique pair $(\mathcal{S}(g, s), \mathcal{S}(g, u))$ of C^{1+} self-renormalisable structures such that $\sigma_{g,s} = \sigma_{\mathcal{S}(g,s),s}$ and $\sigma_{g,u} = \sigma_{\mathcal{S}(g,u),u}$. By Lemma 8.5, $g \in \mathcal{T}(\nu, \delta_s, P_s, \delta_u, P_u)$ if, and only if, $(\mathcal{S}(g, s), \mathcal{S}(g, u)) \in \mathcal{D}^s(\nu, \delta_s, P_s) \times \mathcal{D}^u(\nu, \delta_u, P_u)$. By Lemma 8.3 (ii), for ι equal to s and u , $\mathcal{S}(g, \iota) \in \mathcal{D}^\iota(\nu, \delta_\iota, P_\iota)$ if, and only if, $\mathcal{S}(g, \iota)$ is contained in the (δ_ι, P_ι) -bounded solenoid equivalence class of ν which ends the proof.

Proof of (ii). By Lemma 8.5, $g_1 \in \mathcal{T}(\nu_1, \delta_s, P_s, \delta_u, P_u)$ and $g_2 \in \mathcal{T}(\nu_2, \delta_s, P_s, \delta_u, P_u)$ if, and only if, for ι equal to s and u , $\mathcal{S}(g_1, \iota) \in \mathcal{D}^\iota(\nu_1, \delta_\iota, P_\iota)$ and $\mathcal{S}(g_2, \iota) \in \mathcal{D}^\iota(\nu_2, \delta_\iota, P_\iota)$. By Lemma 9.1,

$\mathcal{S}(g_1, \iota)$ and $\mathcal{S}(g_2, \iota)$ are Lipschitz conjugate if, and only if, $\nu_1 = \nu_2$. Since, by Lemma 9.2, g_1 and g_2 are Lipschitz conjugate if, and only if, for ι equal to s and u , $\mathcal{S}(g_1, \iota)$ and $\mathcal{S}(g_2, \iota)$ are Lipschitz conjugate, we get that g_1 and g_2 are Lipschitz conjugate if, and only if, $\nu_1 = \nu_2$. $\square$

9.2. Extending the result of De la Llave, Marco and Moriyon. Let $\mathcal{P}$ be the set of all periodic points in Λ under f . Let $p(x)$ be the (smallest) period of the periodic point $x \in \mathcal{P}$. For every $x \in \mathcal{P}$ and $\iota \in \{s, u\}$, let $j : J \rightarrow \mathbb{R}$ be a chart in $\mathcal{A}(g, \rho_g)$ such that $x \in J$. The *eigenvalue* $\lambda_{g, \iota}^x(x)$ of x is the derivative of the map $j^{-1}f^p j$ at $j(x)$.

For $\iota \in \{s, u\}$, by construction of the train-tracks, $\mathcal{P}^\iota = \pi_{\mathbf{B}^\iota}(\mathcal{P})$ is the set of all periodic points in $\mathbf{B}^\iota$ under the Markov map f_ι . Furthermore, $\pi_{\mathbf{B}^\iota}|_{\mathcal{P}}$ is an injection and the periodic points $x \in \Lambda$ and $\pi_{\mathbf{B}^\iota}(x) \in \mathbf{B}^\iota$ have the same period $p(x) = p(\pi_{\mathbf{B}^\iota}(x))$. Let us denote $\pi_{\mathbf{B}^\iota}(x)$ by x_ι . Let $\mathcal{S}_\iota$ be a C^{1+} self-renormalisable structure. Let $j : J \rightarrow \mathbb{R}$ be a train-track chart of $\mathcal{S}_\iota$ such that $x_\iota \in J$. The *eigenvalue* $\lambda_{\mathcal{S}_\iota}(x_\iota)$ of x_ι is the derivative of the map $j \circ \tau_\iota^{p(x_\iota)} \circ j^{-1}$ at $j(x_\iota)$, where τ_ι is the Markov map on the train-track $\mathbf{B}^\iota$.

For every $x \in \mathcal{P}$, every $\iota \in \{s, u\}$ and every $n \geq 0$, let $I_n^\iota(x)$ be an ι -leaf $(np(x) + 1)$ -cylinder segment such that $x \in I_n^\iota(x)$ and $f_\iota^{p(x)} I_{n+1}^\iota(x) = I_n^\iota(x)$.

Lemma 9.4. *For $\iota \in \{s, u\}$, let $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota, P_\iota)$ be a C^{1+} ι self-renormalisable structure. For every $x \in \mathcal{P}$,*

$$\begin{aligned} (9.1) \quad \lambda_{\mathcal{S}_\iota}(x_\iota) &= r_{\mathcal{S}_\iota}(I_0^\iota(x) : I_1^\iota(x)) \\ (9.2) \quad &= \rho_{\nu, \iota}(I_0^\iota : I_1^\iota)^{-1/\delta_\iota} e^{-p(x)P_\iota/\delta_\iota} \\ (9.3) \quad &= \rho_{\nu, \iota'}(I_0^{\iota'} : I_1^{\iota'})^{-1/\delta_\iota} e^{-p(x)P_\iota/\delta_\iota}, \end{aligned}$$

where $r_{\mathcal{S}_\iota}$ is the ι -ratio function of $\mathcal{S}_\iota$, $\rho_{\nu, \iota}$ is the ι -measure ratio function of the Gibbs measure ν , and $\rho_{\nu, \iota'}$ is the ι' -measure ratio function of the Gibbs measure ν .

Proof. For every $x \in \mathcal{P}$, let us denote by p the period $p(x)$ of x , and let us denote by I_n^ι the interval $I_n^\iota(x)$. We note that the p -mother $m^p I_{n+1}^\iota$ of I_{n+1}^ι is I_n^ι , and so $f_\iota^p I_{n+1}^\iota = m^p I_{n+1}^\iota$. By (4.1),

$$r_{\mathcal{S}_\iota}(I_0^\iota : I_1^\iota) = \lim_{n \rightarrow \infty} \frac{|I_n^\iota|}{|I_{n+1}^\iota|}.$$

Hence,

$$\begin{aligned} \lambda_{\mathcal{S}_\iota}(x_\iota) &= \lim_{n \rightarrow \infty} \frac{|f_\iota^p I_{n+1}^\iota|}{|I_{n+1}^\iota|} \\ &= \lim_{n \rightarrow \infty} \frac{|I_n^\iota|}{|I_{n+1}^\iota|} \\ &= r_{\mathcal{S}_\iota}(I_0^\iota : I_1^\iota) \end{aligned}$$

which proves (9.1). By Theorem 6.8, the ι -measure ratio function $\rho_{\mathcal{S}_\iota, \delta_\iota}$ is the ι -measure ratio function $\rho_{\nu, \iota}$ of the Gibbs measure ν . Hence, by (7.5), we get

$$\begin{aligned} (9.4) \quad r_{\mathcal{S}_\iota}(I_1^\iota : I_0^\iota) &= \prod_{l=0}^{p-1} r_{\mathcal{S}_\iota}(m^l I_1^\iota : m^{l+1} I_1^\iota) \\ &= \prod_{l=0}^{p-1} \left(J_{\mathcal{S}_\iota, \delta_\iota}(\xi_l) \rho_{\nu, \iota}(m^l I_1^\iota : m^{l+1} I_1^\iota)^{1/\delta_\iota} e^{P_\iota/\delta_\iota} \right) \end{aligned}$$

where $\xi_l = f_t^{p-l} m^l I_1^\iota \in \mathbf{B}_o^\iota$. We note that $f_{\iota'} \xi_l = \xi_{l+1}$ and $f_{\iota'} \xi_{p-1} = \xi_0$ in $\mathbf{B}_o^{\iota'}$. Since $J_{\mathcal{S}_\iota, \delta_\iota} = \kappa/(\kappa \circ f_{\iota'})$ for some function κ , we get

$$(9.5) \quad \prod_{l=0}^{p-1} J_{\mathcal{S}_\iota, \delta_\iota}(\xi_l) = \prod_{l=0}^{p-1} \frac{\kappa(\xi_l)}{\kappa(\xi_{l+1})} = 1.$$

Furthermore,

$$(9.6) \quad \prod_{l=0}^{p-1} \rho_{\nu, \iota}(m^l I_1^\iota : m^{l+1} I_1^\iota) = \rho_{\nu, \iota}(I_1^\iota : I_0^\iota).$$

Using (9.5) and (9.6) in (9.4) we obtain that

$$r_{\mathcal{S}_\iota}(I_1^\iota : I_0^\iota) = \rho_{\nu, \iota}(I_1^\iota : I_0^\iota)^{1/\delta_\iota} e^{pP_\iota/\delta_\iota}.$$

Therefore, by (9.1), we have

$$\begin{aligned} \lambda_{\mathcal{S}_\iota}(x_\iota) &= r_{\mathcal{S}_\iota}(I_0^\iota : I_1^\iota) \\ &= \rho_{\nu, \iota}(I_0^\iota : I_1^\iota)^{-1/\delta_\iota} e^{-pP_\iota/\delta_\iota} \end{aligned}$$

which proves (9.2). By Lemma 6.7, there is $0 < \varepsilon < 1$ such that for every $n \geq 0$

$$(9.7) \quad \rho_{\nu, s}(I_{n+1}^s : I_n^s) \in (1 \pm \varepsilon^n) \frac{\mu([I_{n+1}^s, I_1^u])}{\mu([I_n^s, I_1^u])}$$

and

$$(9.8) \quad \rho_{\nu, u}(I_{n+1}^u : I_n^u) \in (1 \pm \varepsilon^n) \frac{\mu([I_1^s, I_{n+1}^u])}{\mu([I_1^s, I_n^u])}.$$

Since $f^{np}([I_1^s, I_{n+1}^u]) = ([I_{n+1}^s, I_1^u])$ and by invariance of μ , we obtain that

$$(9.9) \quad \frac{\mu([I_{n+1}^s, I_1^u])}{\mu([I_n^s, I_1^u])} = \frac{\mu([I_1^s, I_{n+1}^u])}{\mu([I_1^s, I_n^u])}$$

Putting together (9.7), (9.8) and (9.9), we obtain that

$$\rho_{\nu, s}(I_{n+1}^s : I_n^s) \in (1 \pm \varepsilon^n) \rho_{\nu, u}(I_{n+1}^u : I_n^u).$$

Hence, by invariance of $\rho_{\mathcal{S}_{s,s}}$ and $\rho_{\mathcal{S}_{s,u}}$ under f , we obtain

$$\begin{aligned} \rho_{\nu, s}(I_1^s : I_0^s) &= \lim_{n \rightarrow \infty} \rho_{\nu, s}(I_{n+1}^s : I_n^s) \\ &= \lim_{n \rightarrow \infty} \rho_{\nu, u}(I_{n+1}^u : I_n^u) \\ &= \rho_{\nu, u}(I_1^u : I_0^u) \end{aligned}$$

which proves (9.3).

Lemma 9.5. *Let $\mathbf{B}^\iota$ be a (gap or a no-gap) train-track.*

- (i) *The C^{1+} self-renormalisable structures $\mathcal{S}_1 \in \mathcal{D}^\iota(\nu_1, \delta, P)$ and $\mathcal{S}_2 \in \mathcal{D}^\iota(\nu_2, \delta, P)$ have the same eigenvalues for all periodic orbits if, and only if, ν_1 is equal to ν_2 .*
- (ii) *The set of eigenvalues of all periodic orbits of a C^{1+} self-renormalisable structure is a complete invariant of each Lipschitz conjugacy class.*

Statement (ii) of the above lemma for Markov maps is also in [34].

Proof of (i). By Lemma 9.1, the C^{1+} self-renormalisable structures $\mathcal{S}_1 \in \mathcal{D}^\iota(\nu_1, \delta, P)$ and $\mathcal{S}_2 \in \mathcal{D}^\iota(\nu_2, \delta, P)$ are Lipschitz conjugate if, and only if, the Gibbs measures ν_1 and ν_2 are equal. By Lemma 9.4, if the Gibbs measures ν_1 and ν_2 are equal, then $\mathcal{S}_1$ and $\mathcal{S}_2$ have the same eigenvalues for all periodic orbits. Hence, to finish the proof of statement (i), we are going to prove that if the

C^{1+} self-renormalisable structures $\mathcal{S}_1$ and $\mathcal{S}_2$ have the same eigenvalues for all periodic orbits, then the C^{1+} self-renormalisable structures $\mathcal{S}_1$ and $\mathcal{S}_2$ are Lipschitz conjugate.

Without loss of generality, let us assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are unstable C^{1+} self-renormalisable structures. For $j \in \{1, 2\}$, the (restricted) u -scaling function $z_{u,j} : \Theta^u \rightarrow \mathbb{R}^+$ of $\mathcal{S}$ is well-defined by (see Section 4.6)

$$z_{u,j}(w_0 w_1 \dots) = \lim_{n \rightarrow \infty} \frac{|\pi_{\mathbf{B}^s} \circ f^{n+1} \circ \pi_{\mathbf{B}^u}^{-1} \circ i_u(w_0 w_1 \dots)|_{k_n}}{|\pi_{\mathbf{B}^s} \circ f^n \circ \pi_{\mathbf{B}^u}^{-1} \circ i_u(w_1 w_2 \dots)|_{k_n}},$$

where k_n is a train-track chart in a C^{1+} self-renormalisable atlas $\mathcal{B}_j$ determined by $\mathcal{S}_j$ such that the domain of the chart k_n contains $\pi_{\mathbf{B}^s} \circ f^n \circ \pi_{\mathbf{B}^u}^{-1} \circ i_u(w_1 w_2 \dots)$. For every stable-leaf $(i+1)$ -cylinder J , let $w(J) \in \Theta^u$ be such that $i_u(w(J)) = \pi_{\mathbf{B}^u}(J)$. Hence, for every $l \in \{0, \dots, i-1\}$ we have that

$$\pi_{\mathbf{B}^u}^{-1} \circ i_u(f_u^l w(J)) = f^{-i+l}(m^l J),$$

where $f^{-i+l}(m^l J)$ are stable-leaf primary cylinders. By construction of the (restricted) u -scaling function $z_{u,j}$ and of the u -scaling function $s_{u,j}$ of $\mathcal{S}_j$, we have that

$$(9.10) \quad s_{u,j}(J : m^i J) = \prod_{l=0}^{i-1} z_{u,j}(f_u^l(w(J))).$$

Let $\mathcal{P}_{\Theta^u}$ be the set of all periodic point under the shift. For every $w = w_0 w_1 \dots \in \mathcal{P}_{\Theta^u}$ let $p(w)$ be the smallest period of w . By construction of the train-tracks, for every w , there is a unique periodic point $x(w) \in \Lambda$ with period $p(w)$ with respect to the map f such that $i_u(w) = \pi_{\mathbf{B}^u} x(w)$. Furthermore, there is a unique periodic point $\pi_s x(w) \in \mathbf{B}^s$ with period $p(w)$ for the Markov map. By (9.10), for every $w \in \mathcal{P}_{\Theta^u}$ we have that

$$(9.11) \quad \prod_{i=0}^{p(w)-1} z_{u,j}(f_u^i(w)) = \lambda_{\mathcal{S}_j}(\pi_{\mathbf{B}^s} x(w)).$$

Since the C^{1+} self-renormalisable structures $\mathcal{S}_1$ and $\mathcal{S}_2$ have the same eigenvalues for all periodic orbits, by (9.11), we have that

$$(9.12) \quad \prod_{i=0}^{p(w)-1} \frac{z_{u,1}(f_u^i(w))}{z_{u,2}(f_u^i(w))} = 1,$$

for every $w \in \mathcal{P}_{\Theta^u}$. From Livšic's theorem (e.g. see [12]) we get that

$$(9.13) \quad \frac{z_{u,1}(w)}{z_{u,2}(w)} = \frac{\kappa(w)}{\kappa \circ f_u(w)}$$

where $\kappa : \Theta^u \rightarrow \mathbb{R}^+$ is a positive Hölder continuous function. By (9.10) and (9.13), for every stable-leaf $(i+1)$ -cylinder J we obtain that

$$(9.14) \quad \begin{aligned} \frac{s_{u,1}(J : m^i J)}{s_{u,2}(J : m^i J)} &= \prod_{l=0}^{i-1} \frac{z_{u,1}(f_u^l(w(J)))}{z_{u,2}(f_u^l(w(J)))} \\ &= \frac{\kappa(w)}{\kappa \circ f_u^i(w)}. \end{aligned}$$

Since κ is bounded away from zero and infinity, there is $C > 1$ such that for all $w \in \Theta^u$ and $i \geq 1$ we have that

$$(9.15) \quad C^{-1} < \frac{\kappa(w)}{\kappa \circ \tau_u^i(w)} < C.$$

Putting together (9.14) and (9.15), we obtain that

$$\frac{s_{u,1}(J : m^i J)}{s_{u,2}(J : m^i J)}.$$

Therefore, the ι -solenoid functions $\sigma_{u,1} : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ and $\sigma_{u,2} : \mathbf{S}^\iota \rightarrow \mathbb{R}^+$ corresponding to the C^{1+} self-renormalisable structures $\mathcal{S}_1$ and $\mathcal{S}_2$ are in the same bounded equivalence class (see Definition 3.2). Hence, by Lemma 9.1, the C^{1+} self-renormalisable structures $\mathcal{S}_1$ and $\mathcal{S}_2$ are Lipschitz conjugate.

Proof of (ii). Statement (ii) follows from putting together Lemma 9.1 and Statement (i) of this lemma with $P = 0$. $\square$

Theorem 9.6. (i) If $g \in \mathcal{T}(f, \Lambda)$ then $\lambda_{g,s}(x) = \lambda_{\mathcal{S}_s}(x_s)^{-1}$ and $\lambda_{g,u}(x) = \lambda_{\mathcal{S}_s}(x_u)$ where, for $\iota \in \{s, u\}$, $\lambda_{g,\iota}(x)$ is the eigenvalue of the C^{1+} hyperbolic diffeomorphism g and $\lambda_{\mathcal{S}_\iota}$ is the eigenvalue of the C^{1+} self-renormalisable structure $\mathcal{S}_\iota = \mathcal{S}(g, \iota)$.
(ii) The set of stable and unstable eigenvalues of all periodic orbits of a C^{1+} hyperbolic diffeomorphism $g \in \mathcal{T}(f, \Lambda)$ is a complete invariant of each Lipschitz conjugacy class.

Proof of (i). By construction of the train-track atlas $\mathcal{B}^\iota(g, \rho_g)$ from the lamination atlas $\mathcal{A}^\iota(g, \rho_g)$ in Section 4.2, if $\lambda_{g,\iota}(x)$ is the eigenvalue of $x \in \mathcal{P}$ then the eigenvalue of $x_\iota \in \mathcal{P}_\iota$ is either $\lambda_{g,\iota}(x)$ if $\iota = u$, or $\lambda_{g,\iota}^{-1}$ if $\iota = s$.

Proof of (ii). By Lemma 9.5, the set of eigenvalues of all periodic orbits of a C^{1+} self-renormalisable structure is a complete invariant of each Lipschitz conjugacy class of C^{1+} self-renormalisable structures. Hence, using Lemma 9.2, we get that the set of stable and unstable eigenvalues of all periodic orbits of a C^{1+} hyperbolic diffeomorphism g is a complete invariant of each Lipschitz conjugacy class. $\square$

9.3. Extending the eigenvalue formula of A. N. Livšic and Ja. G. Sinai. In Lemma 9.4, we give an explicit formula for the stable and unstable eigenvalues of all periodic points of the C^{1+} hyperbolic diffeomorphisms in terms of measure ratio functions. We use this formula, in Theorem 9.7 below, to extend the eigenvalue formula of A. N. Livšic and Ja. G. Sinai for Anosov diffeomorphisms to C^{1+} hyperbolic diffeomorphisms.

Theorem 9.7. A C^{1+} hyperbolic diffeomorphism $g \in \mathcal{T}(f, \Lambda)$ has a natural geometric measure $\mu_{g,\delta_s,\delta_u}$ with pressures $P_s = P_s(g, \delta_s, \delta_u)$ and $P_u = P_u(g, \delta_s, \delta_u)$ if, and only if, for all $x \in \Lambda$

$$(9.16) \quad \lambda_{g,s}(x_s)^{-\delta_s} e^{p(x)P_s} = \lambda_{g,u}(x_u)^{\delta_u} e^{p(x)P_u}.$$

Proof. By Theorem 6.1, the C^{1+} self-renormalisable structures $\mathcal{S}(g, s)$ and $\mathcal{S}(g, u)$ are C^{1+} realisations of Gibbs measures $\nu_1 = \nu_{\mathcal{S}(g,s),\delta_s}$ and $\nu_2 = \nu_{\mathcal{S}(g,u),\delta_u}$. By Lemma 9.4 and the statement (i) of Theorem 9.6, for all $x \in \mathcal{P}$ we have

$$(9.17) \quad \begin{aligned} \lambda_{g,u}(x_u) &= \lambda_{\mathcal{S}(g,u)}(x_u) \\ &= \rho_{\nu_2,u}(I_0^u : I_1^u)^{-1/\delta_u} e^{-p(x)P_u/\delta_u} \end{aligned}$$

and

$$(9.18) \quad \begin{aligned} \lambda_{g,s}(x_s) &= \lambda_{\mathcal{S}(g,s)}(x_s)^{-1} \\ &= \rho_{\nu_1,u}(I_0^u : I_1^u)^{1/\delta_s} e^{p(x)P_s/\delta_s}. \end{aligned}$$

Let us prove that if the C^{1+} hyperbolic diffeomorphism g has a natural geometric measure then (9.16) holds. Hence, by Lemma 8.4, the Gibbs measures ν_1 and ν_2 are equal. By (9.17), we have

$$(9.19) \quad \begin{aligned} \rho_{\nu_1,u}(I_0^u : I_1^u) &= \rho_{\nu_2,u}(I_0^u : I_1^u) \\ &= \lambda_{g,u}(x_\iota)^{-\delta_u} e^{-p(x)P_u}. \end{aligned}$$

By (9.18), we obtain that

$$(9.20) \quad \rho_{\nu_1, u}(I_0^u : I_1^u) = \lambda_{g, s}(x_s)^{\delta_s} e^{-p(x)P_s} .$$

Putting together (9.19) and (9.20), we obtain that

$$\lambda_{g, s}(x_s)^{-\delta_s} e^{p(x)P_s} = \lambda_{g, u}(x_u)^{\delta_u} e^{p(x)P_u} ,$$

and so (9.16) holds. Conversely, let us prove that if (9.16) holds then the C^{1+} hyperbolic diffeomorphism g has a natural geometric measure. Putting together (9.16) and (9.18), we obtain that

$$\lambda_{g, u}(x_u) = \rho_{\nu_1, u}(I_0^u : I_1^u)^{-1/\delta_u} e^{-nP_u/\delta_u} .$$

Hence, the Gibbs measure ν_1 determines the same set of eigenvalues for all periodic orbits of self-renormalisable structures in $\mathbf{B}^u$ as the Gibbs measure ν_2 . Therefore, by Lemma 9.5, $\nu_1 = \nu_2$ and consequently, by Lemma 8.4, the C^{1+} hyperbolic diffeomorphism g has a natural geometric measure.

10. INVARIANT HAUSDORFF MEASURES

In this section, we present the proofs of all theorems stated in the Introduction. Let $\mathcal{S}_\iota$ be a C^{1+} ι self-renormalisable structure. By Lemma 6.3, a natural geometric measure $\mu_{\mathcal{S}_\iota, \delta_\iota}$ with pressure $P(\mathcal{S}_\iota, \delta_\iota) = 0$ is an invariant measure absolutely continuous with respect to the Hausdorff measure of $\mathbf{B}^\iota$ and δ_ι is the Hausdorff dimension of $\mathbf{B}^\iota$ with respect to the charts of $\mathcal{S}_\iota$. Let us denote $\mathcal{D}^\iota(\nu, \delta_\iota, 0)$ and $\mathcal{JG}^\iota(\nu, \delta_\iota, 0)$ respectively by $\mathcal{D}^\iota(\nu, \delta_\iota)$ and $\mathcal{JG}^\iota(\nu, \delta_\iota)$. By Theorem 6.1, for every C^{1+} ι self-renormalisable structure $\mathcal{S}_\iota$ there is a unique Gibbs measure $\nu_{\mathcal{S}_\iota}$ such that $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota)$. Using Lemma 8.5, we obtain that the sets $[\nu] \subset \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$ defined in the introduction are equal to the sets $\mathcal{T}(\nu, \delta_s, 0, \delta_u, 0)$ (see Definition 8.3).

Theorem 1.1 follows from Theorem 9.7, Theorem 1.8 follows from Theorem 8.7, Theorem 1.10 follows from Lemma 4.2, and Theorem 1.11 follows from Lemma 8.5.

Proof of Theorem 1.2. Proof of statement (i). By Lemma 9.3 (ii), the sets $[\nu] \subset \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$ are Lipschitz conjugacy classes in $\mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$, and the map $\nu \rightarrow \mathcal{T}(\nu, \delta_s, \delta_u)$ is injective. If $g \in \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$ then g has a natural geometric measure $\mu_{g, \delta_s, \delta_u}$ with pressures $P_s(g, \delta_s, \delta_u)$ and $P_u(g, \delta_s, \delta_u)$ equal to zero. By Lemma 8.4, there is a Gibbs measure $\nu = \nu_{g, \delta_s, \delta_u}$ on Θ such that $i_*\nu = \mu_{g, \delta_s, \delta_u}$ and so $g \in [\nu] \subset \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$. Hence, the map $\nu \rightarrow \mathcal{T}(\nu, \delta_s, \delta_u)$ is surjective into the Lipschitz conjugacy classes in $\mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$.

Proof of statement (ii). By Theorem 9.6 (ii), the set of stable and unstable eigenvalues of all periodic orbits of a C^{1+} hyperbolic diffeomorphisms $g \in \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$ is a complete invariant of each Lipschitz conjugacy class, and by statement (i) of this lemma the sets $\mathcal{T}(\nu, \delta_s, \delta_u)$ are the Lipschitz conjugacy classes in $\mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$. $\square$

Proof of Theorem 1.12. We will separate the proof in three parts. In part (i), we prove that if $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota)$ then $\sigma_{\nu, \iota}$ satisfies the properties indicated in Theorem 1.12. In part (ii), we prove the converse of part (i). In part (iii), we prove that $\mathcal{D}^\iota(\nu, \delta_\iota) \neq \emptyset$ if, and only if, $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'}) \neq \emptyset$.

Part (i). Let $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota)$. By Lemma 6.8, $\mathcal{S}_\iota$ and δ_ι determine a unique ι -measure ratio function $\rho_{\nu, \iota}$ of the Gibbs measure ν . Hence, the function $\rho_{\nu, \iota} |\text{msol}^{\iota'}$ is the ι -measure solenoid function $\sigma_{\nu, \iota}$ of ν and, by Lemma 5.7, $\sigma_{\nu, \iota}$ satisfies the properties indicated in Theorem 1.12.

Part (ii). Conversely, if ν has an ι -solenoid function $\sigma_{\nu, \iota}$ satisfying the properties indicated in Theorem 1.12, by lemmas 5.4 and 5.7, $\sigma_{\nu, \iota}$ determines an unique ι -measure ratio function $\rho_{\nu, \iota}$ of ν . If $\mathbf{B}^\iota$ is a no-gap train-track, by Lemma 8.1, there is a C^{1+} self-renormalisable structure $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota)$ with $\delta_\iota = 1$. If $\mathbf{B}^\iota$ is a gap train-track then, by Remark 7.10, the set $\mathcal{JG}^\iota(\nu, \delta_\iota)$ is

non-empty (in fact it is an infinite dimensional space). Hence, by Lemma 8.3, the set $\mathcal{D}(\nu, \delta_\iota)$ is also non-empty which ends the proof.

Part (iii). To prove that $\mathcal{D}^\iota(\nu, \delta_\iota) \neq \emptyset$ if, and only if, $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'}) \neq \emptyset$, it is enough to prove one of the implications. Let us prove that if $\mathcal{D}^\iota(\nu, \delta_\iota) \neq \emptyset$ then $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'}) \neq \emptyset$. Let $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota)$. By Lemma 6.8, $\mathcal{S}_\iota$ and δ_ι determine an unique ι -measure ratio function $\rho_{\nu, \iota}$ of the Gibbs measure ν . By Lemma 6.9, the ι -measure ratio function $\rho_{\nu, \iota}$ determines an unique dual ι' -measure ratio function $\rho_{\nu, \iota'}$ of ν . Hence, the function $\rho_{\nu, \iota'}|_{\text{msol}^{\iota'}}$ is the ι -measure solenoid function $\sigma_{\nu, \iota'}$ of ν and, by Lemma 5.7, $\sigma_{\nu, \iota'}$ satisfies the properties indicated in Theorem 1.12. Now the proof follows as in part (ii), with ι changed by ι' , which shows that $\sigma_{\nu, \iota'}$ determines a non-empty set $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'})$. $\square$

Proof of Theorem 1.4 and Theorem 1.5. Proof that statement (i) implies statements (ii) and (iii). If $g \in [\nu] \subset \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$, by Lemma 8.5, the sets $\mathcal{D}^s(\nu, \delta_s)$ and $\mathcal{D}^u(\nu, \delta_u)$ are both non-empty. Hence, by Theorem 1.12, the stable measure solenoid function of the Gibbs measure ν satisfies (ii) and the unstable measure solenoid function of the Gibbs measure ν satisfies (iii).

Proof that statement (ii) implies statement (i), and that statement (iii) implies statement (i). By Theorem 1.12, the properties of the ι -solenoid function $\sigma_{\nu, \iota}$ indicated in this theorem imply that $\mathcal{D}^\iota(\nu, \delta_\iota) \neq \emptyset$. Again, by Theorem 1.12 and $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'}) \neq \emptyset$. Hence, by Lemma 8.5, the set $[\nu] \subset \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$ is non-empty. Therefore, every $g \in \mathcal{T}(\nu, \delta_s, \delta_u)$ is a C^{1+} -Hausdorff realisation of ν which ends the proof. $\square$

Proof of Theorem 1.6. Let ν be a Gibbs measure. By Theorem 1.12, the set $\mathcal{D}^s(\nu, \delta_s)$ and $\mathcal{D}^u(\nu, \delta_u)$ are both non-empty. Hence, by Lemma 8.5, the set $\mathcal{T}(\nu, \delta_s, \delta_u)$ is also non-empty. Therefore, every $g \in \mathcal{T}(\nu, \delta_s, \delta_u)$ is a C^{1+} -Hausdorff realisation of ν which ends the proof. $\square$

10.1. Moduli space SOL^ι . Recall the definition of the set SOL^ι given in Section 5.5. By Theorem 10.1, below, the set of all ι -measure solenoid functions σ_ν with the properties indicated in Theorem 1.12 determine an infinite dimensional metric space SOL^ι which gives a nice parametrization of all Lipschitz conjugacy classes $\mathcal{D}^\iota(\nu, \delta)$ of C^{1+} self-renormalisable structures $\mathcal{S}_\iota$ with a given Hausdorff dimension δ .

Theorem 1.7 follows from Theorem 10.1.

Theorem 10.1. *If $\mathbf{B}^\iota$ is a gap train-track assume $0 < \delta_\iota < 1$ and if $\mathbf{B}^\iota$ is a no-gap train-track assume $\delta_\iota = 1$.*

- (i) *The map $\mathcal{S} \rightarrow \rho_{\mathcal{S}, \delta_\iota}$ induces an one-to-one correspondence between the sets $\mathcal{D}^\iota(\nu, \delta_\iota)$ and the elements of SOL^ι .*
- (ii) *The map $g \rightarrow \rho_{\mathcal{S}(g, \iota), \delta_\iota}$ induces an one-to-one correspondence between the sets $[\nu]$ contained in $\mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$ and the elements of SOL^ι .*

Proof of Theorem 10.1. Proof of (i). If $\mathcal{S} \in \mathcal{D}^\iota(\nu, \delta_\iota)$ then the Hausdorff dimension of $\mathcal{S}$ is δ_ι , and $\mathcal{S}$ determines an ι -measure ratio function $\rho_{\mathcal{S}, \delta_\iota} = \rho_{\nu, \iota}$ which does not depend upon $\mathcal{S} \in \mathcal{D}^\iota(\nu, \delta_\iota)$. By Lemma 5.7, $\rho_{\nu, \iota}|_{\text{Msol}^\iota}$ is an element of SOL^ι . Hence, the map $\mathcal{S} \rightarrow \rho_{\mathcal{S}, \delta_\iota}$ associates to each set $\mathcal{D}^\iota(\nu, \delta)$ an unique element of SOL^ι . Conversely, let $\hat{\sigma} \in \text{Msol}^\iota$. By Lemma 5.7, $\hat{\sigma}$ determines an unique ι -measure ratio function $\rho_{\hat{\sigma}}$ such that $\rho_{\hat{\sigma}}|_{\text{Msol}^\iota} = \hat{\sigma}$. By Lemma 5.1, the ι -measure ratio function $\rho_{\hat{\sigma}}$ determines a Gibbs measure $\nu_{\hat{\sigma}}$. If $\mathbf{B}^\iota$ is a no-gap train-track then, by Lemma 8.1, $\rho_{\hat{\sigma}}$ determines a non-empty set $\mathcal{D}^\iota(\nu_{\hat{\sigma}}, \delta_\iota)$. If $\mathbf{B}^\iota$ is a gap train-track then, by Remark 7.10, the set $\mathcal{JG}^\iota(\nu, \delta_\iota)$ is non-empty and so, by Lemma 8.3, the set $\mathcal{D}^\iota(\nu_{\hat{\sigma}}, \delta_\iota)$ is also non-empty. Therefore, each element $\hat{\sigma} \in \text{Msol}^\iota$ determines an unique non-empty set $\mathcal{D}^\iota(\nu_{\hat{\sigma}}, \delta_\iota)$ of C^{1+} self-renormalisable structures $\mathcal{S}$ with $\rho_{\mathcal{S}, \delta_\iota}|_{\text{Msol}^\iota} = \hat{\sigma}$.

Proof of (ii). By Lemma 8.4, if $g \in [\nu]$ then $\mathcal{S}(g, \iota) \in \mathcal{D}^\iota(\nu, \delta_\iota)$ and so, by statement (i) of this lemma, $\rho_{\mathcal{S}(g, \iota), \delta_\iota}|_{\text{Msol}^\iota}$ is an element of SOL^ι which does not depend upon $g \in [\nu]$. Conversely,

let $\hat{\sigma} \in \text{Msol}^\iota$. By statement (i) of this lemma, $\hat{\sigma}$ determines an ι -measure ratio function $\rho_{\hat{\sigma}, \iota}$, and a non-empty set $\mathcal{D}^\iota(\nu_{\hat{\sigma}}, \delta_\iota)$. By Lemma 8.4, $\rho_{\hat{\sigma}, \iota}$ determines an unique dual ι' -ratio function $\rho_{\hat{\sigma}, \iota'}$ associated to the Gibbs measure $\nu_{\hat{\sigma}}$. Again, by statement (i) of this lemma, $\rho_{\hat{\sigma}, \iota'}|\text{Msol}^{\iota'}$ determines a non-empty set $\mathcal{D}^{\iota'}(\nu_{\hat{\sigma}}, \delta_{\iota'})$. By Lemma 8.5, the set $\mathcal{D}^s(\nu_{\hat{\sigma}}, \delta_s) \times \mathcal{D}^u(\nu_{\hat{\sigma}}, \delta_u)$ determines an unique non-empty set $[\nu_{\hat{\sigma}}] \subset \mathcal{T}_{f, \Lambda}(\delta_s, \delta_u)$ of hyperbolic diffeomorphisms $g \in [\nu_{\hat{\sigma}}]$ such that $\rho_{\mathcal{S}(g, \iota), \delta_\iota}|\text{Msol}^\iota = \hat{\sigma}$. $\square$

10.2. Moduli space of cocycle-gap pairs. By Lemma 9.1, each set $\mathcal{D}^\iota(\nu, \delta)$ is a Lipschitz conjugacy class. Hence, by Theorem 1.13 proved below, if $\mathbf{B}^\iota$ is a no-gap train-track then the Lipschitz conjugacy class consists of a single C^{1+} self-renormalisable structure. Furthermore, by Lemma 9.5, the set of eigenvalues of all periodic orbits of $\mathcal{S}_\iota$ is a complete invariant of each set $\mathcal{D}^\iota(\nu, \delta)$.

Proof of Theorem 1.13. Statement (i) follows from Lemma 8.1. Now, let us prove statement (ii). By Remark 7.10, the set $\mathcal{JG}^\iota(\nu, \delta)$ is an infinite dimensional space, and by Lemma 8.3, the set $\mathcal{D}^\iota(\nu, \delta)$ is parametrized by the cocycle-gap pairs in $\mathcal{JG}^\iota(\nu, \delta)$ which ends the proof. $\square$

Proof of Theorem 1.14. By Lemma 8.5, if $g \in \mathcal{T}(\nu, \delta_s, \delta_u)$ then $\mathcal{S}_{\iota'}(g) \in \mathcal{D}^{\iota'}(\nu, \delta_{\iota'})$. Conversely, let $\mathcal{S}_{\iota'}$ be a C^{1+} self-renormalisable structure contained in $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'})$. By Lemma 8.5, a pair $(\mathcal{S}_\iota, \mathcal{S}_{\iota'})$ determines a C^{1+} hyperbolic diffeomorphism $g \in \mathcal{T}(\nu, \delta_s, \delta_u)$. if, and only if, $\mathcal{S}_{\iota'} \in \mathcal{D}^{\iota'}(\nu, \delta_{\iota'})$. By Theorem 1.12, the set $\mathcal{D}(\nu, \delta_\iota)$ is non-empty. Noting that $\delta_\iota = 1$, it follows from Theorem 1.13 (ii) that the set $\mathcal{D}^\iota(\nu, \delta_\iota)$ contains only one C^{1+} self-renormalisable structure $\mathcal{S}_\iota$ which finishes the proof. $\square$

10.3. δ_ι -bounded solenoid equivalence class of Gibbs measures. When we speak of a δ_ι -bounded solenoid equivalence class of ν we mean a $(\delta_\iota, 0)$ -bounded solenoid equivalence class of a Gibbs measure ν (see Definition 8.2). In Section 7, we use the cocycle-gap pairs to construct explicitly the solenoid functions in the δ_ι -bounded solenoid equivalence classes of the Gibbs measures ν . By Theorem 1.15 (ii) proved below, given an ι -solenoid function σ_ι there is an unique Gibbs measure ν such that σ_ι belongs to the δ_ι -bounded solenoid equivalence class of ν .

Proof of Theorem 1.15. Statement (i) follows from Lemma 9.3 (i). Statement (ii) follows from Lemma 8.1 if $\mathbf{B}^\iota$ is a no-gap train-track, and from Lemma 8.3 (ii) if $\mathbf{B}^\iota$ is a gap train-track. Statement (iii) follows from statement (ii) and Theorem 1.13. $\square$

Proof of Theorem 1.16. By Theorem 1.15 (ii), the ι -solenoid function σ_ι determines an unique C^{1+} self-renormalisable structure $\mathcal{S}_\iota \in \mathcal{D}^\iota(\nu, \delta_\iota)$. By Theorem 1.12, the set $\mathcal{D}^{\iota'}(\nu, \delta_{\iota'})$ is nonempty. Let $\mathcal{S}_{\iota'} \in \mathcal{D}^{\iota'}(\nu, 1)$. By Theorem 1.15 (ii), the C^{1+} self-renormalisable structure $\mathcal{S}_{\iota'}$ determines an unique ι' -solenoid function $\sigma_{\iota'}$ such that, by Theorem 1.15 (i), the pair $(\sigma_\iota, \sigma_{\iota'})$ determines an unique C^{1+} conjugacy class $\mathcal{T}(\nu, \delta_s, \delta_u)$ of hyperbolic diffeomorphisms $g \in \mathcal{T}(\nu, \delta_s, \delta_u)$ with an invariant measure $\mu = i_*\nu$ absolutely continuous with respect to the Hausdorff measure. $\square$

Acknowledgments

We are grateful to Dennis Sullivan, Mikhail Lyubich, Krerley Oliveira, Carlos Matheus, and Flávio Ferreira for a number of very fruitful and useful discussions on this work and also for their friendship and encouragement. We thank the Calouste Gulbenkian Foundation, PRO-DYN-ESF, POCTI by FCT and Ministério da Ciência e da Tecnologia, and Centro de Matemática da Universidade do Porto for their financial support of A. A. Pinto and the UK Engineering and Physical Sciences Research Council and the EU for their financial support of D. A. Rand. Part of this research was started during a visit by the authors to the IHES and CUNY and continued at the Isaac Newton Institute, IMPA and SUNY. We thank them for their hospitality.

REFERENCES

- [1] V. Arnol'd, Small denominators I: on the mapping of a circle into itself. *Investijia Akad. Nauk Math*, 25, 1, 1961, 21-96. Transl. A.M.S., 2nd series, 46, 213-284.
- [2] R. Bowen, Equilibrium States and the Ergodic Theory of Axiom A Diffeomorphisms. *Lecture Notes in Mathematics*, No. 470, Springer-Verlag, New York, 1975.
- [3] E. Cawley, The Teichmüller space of an Anosov diffeomorphism of T^2 . *Inventiones Mathematicae* **112** (1993) 351-376.
- [4] K. J. Falconer, *The geometry of fractal sets*. CUP 1985.
- [5] E. de Faria, W. de Melo and A. A. Pinto, Global hyperbolicity of renormalization for C^r unimodal mappings. *Annals of Mathematics*. To appear.
- [6] H. Federer, *Geometric Measure Theory*. New York: Springer 1969.
- [7] M. J. Feigenbaum, Presentation functions, fixed points, and a theory of scaling function dynamics. *J. Stat. Phys.* **52** (1988) 527-569.
- [8] M. J. Feigenbaum, Presentation functions, and scaling function theory for circle maps. *Nonlinearity* **1** (1988) 577-602.
- [9] J. Franks, Anosov diffeomorphisms on tori. *Trans. Amer. Math. Soc.* **145** (1969) 117-124.
- [10] M. R. Herman, Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Publ. IHES* **49**, 1979, 5-233.
- [11] M. Hirsch and C. Pugh, Stable manifolds and hyperbolic sets. *Proc. Symp. Pure Math., Amer. Math. Soc.* **14** (1970) 133-164.
- [12] A. Katok, and B. Hasselblatt, *Introduction to The Modern Theory of Dynamical Systems* CUP 1994.
- [13] A. N. Livšic and Ja. G. Sinai, Invariant measures that are compatible with smoothness for transitive C -systems. (Russian) *Dokl. Akad. Nauk SSSR* **207** (1972), 1039-1041.
- [14] R. Llave, J. M. Marco and R. Moriyon, Canonical perturbation theory of Anosov systems and regularity results for the Livšic cohomology equations. *Annals of Math.*, **123** (1986) 537-612.
- [15] R. Llave, Invariants for Smooth conjugacy of hyperbolic dynamical systems II. *Commun. Math. Phys.*, **109** (1987) 369-378.
- [16] R. Mañé, *Ergodic Theory and Differentiable Dynamics*. Springer-Verlag Berlin 1987.
- [17] J. M. Marco and R. Moriyon, Invariants for Smooth conjugacy of hyperbolic dynamical systems I. *Commun. Math. Phys.*, **109** (1987) 681-689.
- [18] J. M. Marco and R. Moriyon, Invariants for Smooth conjugacy of hyperbolic dynamical systems III. *Commun. Math. Phys.*, **112** (1987) 317-333.
- [19] S. Newhouse, On codimension one Anosov diffeomorphisms. *Amer. J. Math.* **92** (1970) 671-762.
- [20] S. Newhouse and J. Palis, Hyperbolic nonwandering sets on two-dimension manifolds. *Dynamical systems* (ed. M Peixoto) (1973).
- [21] A. A. Pinto and D. A. Rand, Classifying C^{1+} structures on hyperbolic fractals: 1 The moduli space of solenoid functions for Markov maps on train-tracks. *Erg. Th. Dyn. Syst.* (1995) 697-734.
- [22] A. A. Pinto and D. A. Rand, Classifying C^{1+} structures on hyperbolic fractals: 2 Embedded trees. *Erg. Th. Dyn. Syst.* (1995) 969-992.
- [23] A. A. Pinto and D. A. Rand, Existence, uniqueness and ratio decomposition for Gibbs states via duality. *Erg. Th. Dyn. Syst.* **21** (2001) 533-544.
- [24] A. A. Pinto and D. A. Rand, Teichmüller spaces and HR structures for hyperbolic surface dynamics. *Erg. Th. Dyn. Syst.* **22** (2002) 1905-1931.
- [25] A. A. Pinto and D. A. Rand, Smoothness of holonomies for codimension 1 hyperbolic dynamics. *Bull. London Math. Soc.* **34** (2002) 341-352.
- [26] A. A. Pinto and D. A. Rand, Rigidity of hyperbolic surface dynamics. *J. Lond. Math. Soc.* **71** (2) (2004) 481-502.
- [27] A. A. Pinto and D. A. Rand, Solenoid functions for hyperbolic sets on surfaces. To be published by *MSRI book series (published by Cambridge University Press)*.
- [28] A. A. Pinto and D. Sullivan, The circle and the solenoid. To be published by *Katok birthday issue of DCDS* 1-37.
- [29] S. Ostlund, D. A. Rand, J. Sethna, and E. Siggia, Universal properties of the transition from quasi-periodicity to chaos. *Physica D*, **8** 1-54 (1983).
- [30] D. A. Rand, Universality and renormalisation in dynamical systems. In *New Directions in Dynamical Systems* (ed. T. Bedford and J. Swift) pp. 1-56, (1988), CUP
- [31] M. Shub, *Global Stability of Dynamical Systems*, Springer-Verlag (1987).
- [32] Ja. G. Sinai, Markov Partitions and C -diffeomorphisms. *Anal. and Appl.* **2** (1968) 70-80.

- [33] D. Sullivan, Differentiable structures on fractal-like sets determined by intrinsic scaling functions on dual Cantor sets. *Proceedings of Symposia in Pure Mathematics* **48** (1988), American Mathematical Society.
- [34] D. Sullivan, Bounds, quadratic differentials, and renormalization conjectures. *AMS Centennial Publications*. Volume 2: Mathematics into the Twenty-first Century (1988 Centennial Symposium, August 8-12), American Mathematical Society, Providence, RI.,1991.
- [35] R. F. Williams, Expanding attractors. *Publ. I.H.E.S.* **43** (1974) 169-203.
- [36] J. C. Yoccoz, Conjugaison différentiable des difféomorphismes du cercle dont le nombre de rotation vérifie une condition diophantienne. *Ann. scient. Éc. Norm. Sup.* 4 série, t, **17** (1984) 333-359.

(A. A. Pinto) FACULDADE DE CIENCIAS, UNIVERSIDADE DO PORTO, 4000 PORTO, PORTUGAL.
E-mail address, A. A. Pinto: `aapinto@fc.up.pt`

(D. A. Rand) MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UK.
E-mail address, D. A. Rand: `dar@maths.warwick.ac.uk`